

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>C12Q 1/68</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 95/09248</b> (43) International Publication Date: <b>6 April 1995 (06.04.95)</b></p>
<p>(21) International Application Number: <b>PCT/US94/10945</b> (22) International Filing Date: <b>27 September 1994 (27.09.94)</b> (30) Priority Data:  <div style="display: flex; justify-content: space-between;"> <div>08/127,420      27 September 1993 (27.09.93)    US</div> <div>08/303,058      8 September 1994 (08.09.94)      US</div> </div>   <p>(60) Parent Applications or Grants (63) Related by Continuation  <div style="display: flex; justify-content: space-between;"> <div>US      08/127,420 (CIP)</div> <div>Filed on      27 September 1993 (27.09.93)</div> </div> <div style="display: flex; justify-content: space-between;"> <div>US      08/303,058 (CIP)</div> <div>Filed on      8 September 1994 (08.09.94)</div> </div>   <p>(71) Applicant (for all designated States except US): <b>ARCH DEVELOPMENT CORP. [US/US]; 1101 E. 58th Street, Chicago, IL 60637 (US).</b> (72) Inventor; and (75) Inventor/Applicant (for US only): <b>DRMANAC, Radoje [YU/US]; Apartment #212, 2622 Burr Ridge Ct., Woodridge, IL 60517 (US).</b></p> </p></p>	<p>(74) Agent: <b>PARKER, David, L.; Arnold, White &amp; Durkee, P.O. Box 4433, Houston, TX 77210 (US).</b>  (81) Designated States: <b>AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).</b>  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: **METHODS AND COMPOSITIONS FOR EFFICIENT NUCLEIC ACID SEQUENCING**

(57) Abstract

Disclosed are novel methods and compositions for rapid and highly efficient nucleic acid sequencing based upon hybridization with two sets of small oligonucleotide probes of known sequences. Extremely large nucleic acid molecules, including chromosomes and non-amplified RNA, may be sequenced without prior cloning or subcloning steps. The methods of the invention also solve various current problems associated with sequencing technology such as, for example, high noise to signal ratios and difficult discrimination, attaching many nucleic acid fragments to a surface, preparing many, longer or more complex probes and labelling more species.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LE	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

Description  
METHODS AND COMPOSITIONS FOR  
EFFICIENT NUCLEIC ACID SEQUENCING

5       The present application is a continuation-in-part of  
co-pending U.S. Patent Application Serial No. 08/303,058,  
filed September 08, 1994; which is a continuation-in-part  
of U.S. Patent Application Serial No. 08/127,420, filed  
September 27, 1993; the entire text and figures of which  
10 disclosures are specifically incorporated herein by  
reference without disclaimer. The U.S. Government owns  
rights in the present invention pursuant to Department of  
Energy grant LDRD 03235 and Contract No. W-31-109-ENG-38  
between the U.S. Department of Energy and The University  
15 of Chicago, representing Argonne National Laboratory.

BACKGROUND OF THE INVENTION

1. Field of the Invention

20

The present invention generally relates to the field  
of molecular biology. The invention particularly  
provides novel methods and compositions to enable highly  
efficient sequencing of nucleic acid molecules. The  
25 methods of the invention are suitable for sequencing long  
nucleic acid molecules, including chromosomes and RNA,  
without cloning or subcloning steps.

2. Description of the Related Art

30

Nucleic acid sequencing forms an integral part of  
scientific progress today. Determining the sequence,  
i.e. the primary structure, of nucleic acid molecules and  
segments is important in regard to individual projects  
35 investigating a range of particular target areas.  
Information gained from sequencing impacts science,  
medicine, agriculture and all areas of biotechnology.

- 2 -

Nucleic acid sequencing is, of course, vital to the human genome project and other large-scale undertakings, the aim of which is to further our understanding of evolution and the function of organisms and to provide an insight into the causes of various disease states.

The utility of nucleic acid sequencing is evident, for example, the Human Genome Project (HGP), a multinational effort devoted to sequencing the entire human genome, is in progress at various centers. However, progress in this area is generally both slow and costly. Nucleic acid sequencing is usually determined on polyacrylamide gels that separate DNA fragments in the range of 1 to 500 bp, differing in length by one nucleotide. The actual determination of the sequence, i.e., the order of the individual A, G, C and T nucleotides may be achieved in two ways. Firstly, using the Maxam and Gilbert method of chemically degrading the DNA fragment at specific nucleotides (Maxam & Gilbert, 1977), or secondly, using the dideoxy chain termination sequencing method described by Sanger and colleagues (Sanger et al., 1977). Both methods are time-consuming and laborious.

More recently, other methods of nucleic acid sequencing have been proposed that do not employ an electrophoresis step, these methods may be collectively termed Sequencing By Hybridization or SBH (Drmanac et al., 1991; Cantor et al., 1992; Drmanac & Crkvenjakov, U.S. Patent 5,202,231). Development of certain of these methods has given rise to new solid support type sequencing tools known as sequencing chips. The utility of SBH in general is evidenced by the fact that U.S. Patents have been granted on this technology. However, although SBH has the potential for increasing the speed with which nucleic acids can be sequenced, all current SBH methods still suffer from several drawbacks.

- 3 -

SBH can be conducted in two basic ways, often referred to as Format 1 and Format 2 (Cantor et al., 1992). In Format 1, oligonucleotides of unknown sequence, generally of about 100-1000 nucleotides in length, are arrayed on a solid support or filter so that the unknown samples themselves are immobilized (Strezoska et al., 1991; Drmanac & Crkvenjakov, U.S. Patent 5,202,231). Replicas of the array are then interrogated by hybridization with sets of labeled probes of about 6 to 8 residues in length. In Format 2, a sequencing chip is formed from an array of oligonucleotides with known sequences of about 6 to 8 residues in length (Southern, WO 89/10977; Khrapko et al., 1991; Southern et al., 1992). The nucleic acids of unknown sequence are then labeled and allowed to hybridize to the immobilized oligos.

Unfortunately, both of these SBH formats have several limitations, particularly the requirement for prior DNA cloning steps. In Format 1, other significant problems include attaching the various nucleic acid pieces to be sequenced to the solid surface support or preparing a large set of longer probes. In Format 2, major problems include labelling the nucleic acids of unknown sequence, high noise to signal ratios that generally result, and the fact that only short sequences can be determined. Further problems of Format 2 include the secondary structure formation that prevents access to some targets and the different conditions that are necessary for probes with different GC contents. Therefore, the art would clearly benefit from a new procedure for nucleic acid sequencing, and particularly, one that avoids the tedious processes of cloning and/or subcloning.

35

- 4 -

SUMMARY OF THE INVENTION

The present invention seeks to overcome these and other drawbacks inherent in the prior art by providing  
5 new methods and compositions for the sequencing of nucleic acids. The novel techniques described herein have been generally termed Format 3 by the inventors and represent marked improvements over the existing Format 1 and Format 2 SBH methods. In the Format 3 sequencing  
10 provided by the invention, nucleic acid sequences are determined by means of hybridization with two sets of small oligonucleotide probes of known sequences. The methods of the invention allow high discriminatory sequencing of extremely large nucleic acid molecules,  
15 including chromosomal material or RNA, without prior cloning, subcloning or amplification. Furthermore, the present methods do not require large numbers of probes, the complex synthesis of longer probes, or the labelling of a complex mixture of nucleic acids segments.

20

To determine the sequence of a nucleic acid according to the methods of the present invention, one would generally identify sequences from the nucleic acid by hybridizing with complementary sequences from two sets  
25 of small oligonucleotide probes (oligos) of defined length and known sequence, which cover most combinations of sequences for that length of probe. One would then analyze the sequences identified to determine stretches of the identified sequences that overlap, and reconstruct  
30 or assemble the complete nucleic acid sequence from such overlapping sequences.

The sequencing methods may be conducted using sequential hybridization with complementary sequences  
35 from the two sets of small oligos. Alternatively, a mode described as "cycling" may be employed, in which the two sets of small oligos are hybridized with the unknown

- 5 -

sequences simultaneously. The term "cycling" is applied as the discriminatory part of the technique comes from then increasing the temperature to "melt" those hybrids that are non-complementary. Such cycling techniques are  
5 commonly employed in other areas of molecular biology, such as PCR, and will be readily understood by those of skill in the art in light when reading the present disclosure.

10 The invention is applicable to sequencing nucleic acid molecules of very long length. As a practical matter, the nucleic acid molecule to be sequenced will generally be fragmented to provide small or intermediate length nucleic acid fragments that may be readily  
15 manipulated. The term nucleic acid fragment, as used herein, most generally means a nucleic acid molecule of between about 10 base pairs (bp) and about 100 bp in length. The most preferred methods of the invention are contemplated to be those in which the nucleic acid  
20 molecule to be sequenced is treated to provide nucleic acid fragments of intermediate length, i.e., of between about 10 bp and about 40 bp. However, it should be stressed that the present invention is not a method of completely sequencing small nucleic acid fragments,  
25 rather it is a method of sequencing nucleic acid molecules *per se*, which involves determining portions of sequence from within the molecule - whether this is done using the whole molecule, or for simplicity, whether this is achieved by first fragmenting the molecule into  
30 smaller sized sections of from about 4 to about 1000 bases.

Sequences from nucleic acid molecules are determined by hybridizing to small oligonucleotide probes of known  
35 sequence. In referring to "small oligonucleotide probes", the term "small" means probes of less than 10 bp in length, and preferably, probes of between about 4 bp

- 6 -

and about 9 bp in length. In one exemplary sequencing embodiment, probes of about 6 bp in length are contemplated to be particularly useful. For the sets of oligos to cover all combinations of sequences for the length of probe chosen, their number will be represented by  $4^F$ , wherein F is the length of the probe. For example, for a 4-mer, the set would contain 256 probes; for a 5-mer, the set would contain 1024 probes; for a 6-mer, 4096 probes; a 7-mer, 16384 probes; and the like.

10 The synthesis of oligos of this length is very routine in the art and may be achieved by automated synthesis.

In the methods of the invention, one set of the small oligonucleotide probes of known sequence, which may be termed the first set, will be attached to a solid support, i.e., immobilized on that support in such a way so that they are available to take part in hybridization reactions. The other set of small oligonucleotide probes of known sequence, which may be termed the second set, will be probes that are in solution and that are labelled with a detectable label. The sets of oligos may include probes of the same or different lengths.

The process of sequential hybridization means that nucleic acid molecules, or fragments, of unknown sequence can be hybridized to the distinct sets of oligonucleotide probes of known sequences at separate times (FIG. 1). The nucleic acid molecules or fragments will generally be denatured, allowing hybridization, and added to the first, immobilized set of probes under discriminating hybridization conditions to ensure that only fragments with complementary sequences hybridize. Fragments with non-complementary sequences are removed and the next round of discriminating hybridization is then conducted by adding the second, labelled set of probes, in solution, to the combination of fragments and probes already formed. Labelled probes that hybridize adjacent



- 7 -

to a fixed probe will remain attached to the support and can be detected, which is not the case when there is space between the fixed and labelled probes (FIG. 1).

5       The process of simultaneous hybridization means that the unknown sequence nucleic acid molecules can be contacted with the distinct sets of oligonucleotide probes of known sequences at the same time. Hybridization will occur under discriminating  
10       hybridization conditions. Fragments with non-complementary sequences are then "melted", i.e., removed by increasing the temperature, and the next round of discriminating hybridization is then conducted, allowing any complementary second probes to hybridize. Labelled  
15       probes that hybridize adjacent to a fixed probe will then be detected in the same manner.

      Nucleic acid sequences that are "complementary" are those that are capable of base-pairing according to the  
20       standard Watson-Crick complementarity rules, and variations of the rules as they apply to modified bases. That is, that the larger purines, or modified purines, will always base pair with the smaller pyrimidines to form only known combinations. These include the standard  
25       pairs of guanine paired with Cytosine (G:C) and Adenine paired with either Thymine (A:T), in the case of DNA, or Adenine paired with Uracil (A:U) in the case of RNA. The use of modified bases, or the so-called Universal Base (M, Nichols et al., 1994) is also contemplated.

30       As used herein, the term "complementary sequences" means nucleic acid sequences that are substantially complementary over their entire length and have very few base mismatches. For example, nucleic acid sequences of  
35       six bases in length may be termed complementary when they hybridize at five out of six positions with only a single mismatch. Naturally, nucleic acid sequences that are

- 8 -

"completely complementary" will be nucleic acid sequences that are entirely complementary throughout their entire length and have no base mismatches.

5       After identifying, by hybridization to the oligos of known sequence, various individual sequences that are part of the nucleic acid fragments, these individual sequences are next analyzed to identify stretches of sequences that overlap. For example, portions of  
10       sequences in which the 5' end is the same as the 3' end of another sequence, or vice versa, are identified. The complete sequence of the nucleic acid molecule or fragment can then be delineated, i.e., it can be reconstructed from the overlapping sequences thus  
15       determined.

      The processes of identifying overlapping sequences and reconstructing the complete sequence will generally be achieved by computational analysis. For example, if a  
20       labelled probe 5'-TTTTTT-3' hybridizes to the spot containing the fixed probe 5'-AAAAAA-3', a 12-mer sequence from within the nucleic acid molecule is defined, namely 5'-AAAAAATTTTTT-3' (SEQ ID NO:1), i.e. the sequence of the two hybridized probes is combined to  
25       reveal a previously unknown sequence. The next question to be answered is which nucleotide follows next after the newly determined 5'-AAAAAATTTTTT-3' (SEQ ID NO:1) sequence. There are four possibilities represented by the fixed probe 5'-AAAAAT-3' and labelled probes  
30       5'-TTTTTA-3' for A; 5'-TTTTTT-3' for T; 5'-TTTTTC-3' for C; and 5'-TTTTTG-3' for G. If, for example, the probe 5'-TTTTTC-3' is positive and the other three are negative, then the assembled sequence is extended to 5'-AAAAAATTTTTTC-3' (SEQ ID NO:2). In the next step, an  
35       algorithm determines which of the labelled probes TTTTCA, TTTTCT, TTTTCC or TTTTCG are positive at the spot containing the fixed probe AAAATT. The process is

- 9 -

repeated until all positive (F + P) oligonucleotide sequences are used or defined as false positives.

The present invention thus provides a very effective way to sequence nucleic acid fragments and molecules of long length. Large nucleic acid molecules, as defined herein, are those molecules that need to be fragmented prior to sequencing. They will generally be of at least about 45 or 50 base pairs (bp) in length, and will most often be longer. In fact, the methods of the invention may be used to sequence nucleic acid molecules with virtually no upper limit on length, so that sequences of about 100 bp, 1 kilobase (kb), 100 kb, 1 megabase (Mb), and 50 Mb or more may be sequenced, up to and including complete chromosomes, such as human chromosomes, which are about 100 Mb in length. Such a large number is well within the scope of the present invention and sequencing this number of bases will require two sets of 8-mers or 9-mers (so that  $F + P \approx 16-18$ ). The nucleic acids to be sequenced may be DNA, such as cDNA, genomic DNA, microdissected chromosome bands, cosmid DNA or YAC inserts, or may be RNA, including mRNA, rRNA, tRNA or snRNA.

The process of determining the sequence of a long nucleic acid molecule involves simply identifying sequences of length F + P from the molecule and combining the sequences using a suitable algorithm. In practical terms, one would most likely first fragment the nucleic acid molecule to be sequenced to produce smaller fragments, such as intermediate length nucleic acid fragments. One would then identify sequences of length F + P by hybridizing, e.g., sequentially hybridizing, the fragments to complementary sequences from the two sets of small oligonucleotide probes of known sequence, as described above. In this manner, the complete nucleic

- 10 -

acid sequence of extremely large molecules can be reconstructed from overlapping sequences of length  $F + P$ .

Whether the nucleic acid to be sequenced is itself  
5 an intermediate length fragment or is first treated to generate such length fragments, the process of identifying sequences from such nucleic acid fragments by hybridizing to two sets of small oligonucleotide probes of known sequence is central to the sequencing methods  
10 disclosed herein. This process generally comprises the following steps:

- (a) contacting the set or array of attached or immobilized oligonucleotide probes with the  
15 nucleic acid fragments under hybridization conditions effective to allow fragments with a complementary sequence to hybridize sufficiently to a probe, thereby forming primary complexes wherein the fragment has both  
20 hybridized and non-hybridized, or "free", sequences;
- (b) contacting the primary complexes with the set of labelled oligonucleotide probes in solution  
25 under hybridization conditions effective to allow probes with complementary sequences to hybridize to a non-hybridized or free fragment sequence, thereby forming secondary complexes wherein the fragment is hybridized to both an  
30 attached (immobilized) probe and a labelled probe;
- (c) removing from the secondary complexes any labelled probes that have not hybridized  
35 adjacent to an attached probe, thereby leaving only adjacent secondary complexes;

- 11 -

- (d) detecting the adjacent secondary complexes by detecting the presence of the label in the labelled probe; and
- 5 (e) identifying oligonucleotide sequences from the nucleic acid fragments in the adjacent secondary complexes by combining or connecting the known sequences of the hybridized attached and labelled probes.

10

The hybridization or 'washing conditions' chosen to conduct either one, or both, of the hybridization steps may be manipulated according to the particular sequencing embodiment chosen. For example, both of the

15 hybridization conditions may be designed to allow oligonucleotide probes to hybridize to a given nucleic acid fragment when they contain complementary sequences, i.e., substantially matching sequences, such as those sequences that hybridize at five out of six positions.

20 The hybridization steps would preferably be conducted using a simple robotic device as is routinely used in current sequencing procedures.

Alternatively, the hybridization conditions may be

25 designed to allow only those oligonucleotide probes and fragments that have completely complementary sequences to hybridize. These more discriminating or 'stringent' conditions may be used for both distinct steps of a sequential hybridization process or for either step

30 alone. In such cases, the oligonucleotide probes, whether immobilized or labelled probes, would only be allowed to hybridize to a given nucleic acid fragment when they shared completely complementary sequences with the fragment.

35

The hybridization conditions chosen will generally dictate the degree of complexity required to analyze the

- 12 -

data obtained. Equally, the computer programs available to analyze any data generated may dictate the hybridization conditions that must be employed in a given laboratory. For example, in the most discriminating  
5 process, both hybridization steps would be conducted under conditions that allow only oligos and fragments with completely complementary sequences to hybridize. As there will be no mismatched bases, this method involves the least complex computational analyses and, for this  
10 reason, it is the currently preferred method for practicing the invention. However, the use of less discriminating conditions for one or both hybridization steps also falls within the scope of the present invention.

15

Suitable hybridization conditions for use in either or both steps may be routinely determined by optimization procedures or 'pilot studies'. Various types of pilot studies are routinely conducted by those skilled in the  
20 art of nucleic acid sequencing in establishing working procedures and in adapting a procedure for use in a given laboratory. For example, conditions such as the temperature; the concentration of each of the components; the length of time of the steps; the buffers used and  
25 their pH and ionic strength may be varied and thereby optimized.

In preferred embodiments, the nucleic acid sequencing method of the invention involves a  
30 discriminating step to select for secondary hybridization complexes that include immediately adjacent immobilized and labelled probes, as distinct from those that are not immediately adjacent and are separated by one, two or more bases. A variety of processes are available for  
35 removing labelled probes that are not hybridized immediately adjacent to an attached probe, i.e., not

- 13 -

hybridized back to back, each of which leaves only the immediately adjacent secondary complexes.

Such discriminatory processes may rely solely on  
5 washing steps of controlled stringency wherein the hybridization conditions employed are designed so that immediately adjacently probes remain hybridized due to the increased stability afforded by the stacking interactions of the adjacent nucleotides. Again, washing  
10 conditions such as temperature, concentration, time, buffers, pH, ionic strength and the like, may be varied to optimize the removal of labelled probes that are not immediately adjacent.

15 In preferred embodiments the immediately adjacent immobilized and labelled probes would be ligated, i.e., covalently joined, prior to performing washing steps to remove any non-ligated probes. Ligation may be achieved by treating with a solution containing a chemical  
20 ligating agent, such as, e.g., water-soluble carbodiimide or cyanogen bromide. More preferably, a ligase enzyme, such as T<sub>4</sub> DNA ligase from T<sub>4</sub> bacteriophage, which is commercially available from many sources (e.g., Biolabs), may be employed. In any event, one would then be able to  
25 remove non-immediately adjacent labelled probes by more stringent washing conditions that cannot affect the covalently connected labeled and fixed probes.

The remaining adjacent secondary complexes would be  
30 detected by observing the location of the label from the labelled probes present within the complexes. The oligonucleotide probes may be labeled with a chemically-detectable label, such as fluorescent dyes, or adequately modified to be detected by a chemiluminescent developing  
35 procedure, or radioactive labels such as <sup>35</sup>S, <sup>3</sup>H, <sup>32</sup>P or <sup>33</sup>P, with <sup>33</sup>P currently being preferred. Probes may also

- 14 -

be labeled with non-radioactive isotopes and detected by mass spectrometry.

Currently, the most preferred method contemplated  
5 for practicing the present invention involves performing the hybridization steps under conditions designed to allow only those oligonucleotide probes and fragments that have completely complementary sequences to hybridize and that allow only those probes that are immediately  
10 adjacent to remain hybridized. This method subsequently requires the least complex computational analysis.

Where the nucleic acid molecule of unknown sequence is longer than about 45 or 50 bp, one effective method  
15 for determining its sequence generally involves treating the molecule to generate nucleic acid fragments of intermediate length, and determining sequences from the fragments. The nucleic acid molecule, whether it be DNA or RNA may be fragmented by any one of a variety of  
20 methods including, for example, cutting by restriction enzyme digestion, shearing by physical means such as ultrasound treatment, by NaOH treatment or by low pressure shearing.

25 In certain embodiments, e.g., involving small oligonucleotide probes between about 4 bp and about 9 bp in length, one may aim to produce nucleic acid fragments of between about 10 bp and about 40 bp in length. Naturally, longer length probes would generally be used  
30 in conjunction with sequencing longer length nucleic acid fragment, and vice versa. In certain preferred embodiments, the small oligonucleotide probes used will be about 6 bp in length and the nucleic acid fragments to be sequenced will generally be about 20 bp in length. If  
35 desired, fragments may be separated by size to obtain those of an appropriate length, e.g., fragments may be



- 15 -

run on a gel, such as an agarose gel, and those with approximately the desired length may be excised.

5 The method for determining the sequence of a nucleic acid molecule may also be exemplified using the following terms. Initially one would randomly fragment an amount of the nucleic acid to be sequenced to provide a mixture of nucleic acid fragments of length T. One would prepare an array of immobilized oligonucleotide probes of known  
10 sequences and length F and a set of labelled oligonucleotide probes in solution of known sequences and length P, wherein  $F + P \leq T$  and, preferably, wherein  $T \approx 3F$ .

15 One would then contact the array of immobilized oligonucleotide probes with the mixture nucleic acid fragments under hybridization conditions effective to allow the formation of primary complexes with hybridized, complementary sequences of length F and non-hybridized  
20 fragment sequences of length  $T - F$ . Preferably, the hybridized sequences of length F would contain only completely complementary sequences.

The primary complexes would then be contacted with  
25 the set of labelled oligonucleotide probes under hybridization conditions effective to allow the formation of secondary complexes with hybridized, complementary sequences of length F and adjacent hybridized, complementary sequences of length P. In preferred  
30 embodiments, only those labelled probes with completely complementary sequences would be allowed to hybridize and only those probes that hybridize immediately adjacent to an immobilized probe would be allowed to remain hybridized. In the most preferred embodiments, the  
35 adjacent immobilized and labelled oligonucleotide probes would also be ligated at this stage.

- 16 -

Next one would detect the secondary complexes by detecting the presence of the label and identify sequences of length  $F + P$  from the nucleic acid fragments in the secondary complexes by combining the known  
5 sequences of the hybridized immobilized and labelled probes. Stretches of the sequences of length  $F + P$  that overlap would then be identified, thereby allowing the complete nucleic acid sequence of the molecule to be reconstructed or assembled from the overlapping sequences  
10 determined.

In the methods of the invention, the oligonucleotides of the first set may be attached to a solid support, i.e. immobilized, by any of the methods  
15 known to those of skill in the art. For example, attachment may be via addressable laser-activated photodeprotection (Fodor et al., 1991; Pease et al., 1994). One generally preferred method is to attach the oligos through the phosphate group using reagents such as  
20 nucleoside phosphoramidite or nucleoside hydrogen phosphate, as described by Southern & Maskos (PCT Patent Application WO 90/03382, incorporated herein by reference), and using glass, nylon or teflon supports. Another preferred method is that of light-generated  
25 synthesis described by Pease et al. (1994; incorporated herein by reference). One may also purchase support bound oligonucleotide arrays, for example, as have been offered for sale by Affymetrix and Beckman.

30 The immobilized oligonucleotides may be formed into an array comprising all probes or subsets of probes of a given length (preferably about 4 to 10 bases), and more preferably, into multiple arrays of immobilized oligonucleotides arranged to form a so-called "sequencing  
35 chip". One example of a chip is that where hydrophobic segments are used to create distinct spatial areas. The sequencing chips may be designed for different

- 17 -

applications like mapping, partial sequencing, sequencing of targeted regions for diagnostic purposes, mRNA sequencing and large scale genome sequencing. For each application, a specific chip may be designed with  
5 different sized probes or with an incomplete set of probes.

In one exemplary embodiment, both sets of oligonucleotide probes would be probes of six bases in  
10 length, i.e., 6-mers. In this instance, each set of oligos contains 4096 distinct probes. The first set probes is preferably fixed in an array on a microchip, most conveniently arranged in 64 rows and 64 columns. The second set of 4096 oligos would be labeled with a  
15 detectable label and dispensed into a set of distinct tubes. In this example, 4096 of the chips would be combined in a large array, or several arrays. After hybridizing the nucleic acid fragments, a small amount of the labeled oligonucleotides would be added to each  
20 microchip for the second hybridization step, only one of each of the 4096 nucleotides would be added to each microchip.

Further embodiments of the invention include kits  
25 for use in nucleic acid sequencing. Such kits will generally comprise a solid support having attached an array of oligonucleotide probes of known sequences, as shown in FIG. 2A, FIG. 2B and FIG. 2C, wherein the oligonucleotides are capable of taking part in  
30 hybridization reactions, and a set of containers comprising solutions of labelled oligonucleotide probes of known sequences. Arrangements such as those shown in FIG. 4 are also contemplated. This depicts the use of the Universal Base, either as an attachment method, or at  
35 the terminus to give an added dimension to the hybridization of fragments.

- 18 -

In the kits, the attached oligonucleotide probes and those in solution may be between about 4 bp and about 9 bp in length, with ones of about 6 bp in length being preferred. The oligos may be labelled with chemically-detectable or radioactive labels, with  $^{32}\text{P}$ -labelled probes being generally preferred, and  $^{33}\text{P}$ -labelled probes being even more preferred. The kits may also comprise a chemical or other ligating agent, such as a DNA ligase enzyme. A variety of other additional compositions and materials may be included in the kits, such as 96-tip or 96-pin devices, buffers, reagents for cutting long nucleic acid molecules and tools for the size selection of DNA fragments. The kits may even include labelled RNA probes so that the probes may be removed by RNAase treatment and the sequencing chips re-used.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Basic steps in the hybridization process. Step 1: The unlabelled target DNA to be sequenced (T) is hybridized under discriminative conditions to an array of attached oligonucleotide probes. Spots with probe Fx and Fy are depicted. Complementary sequences for Fx and Fy are at different positions of T. Step 2: Labeled probes, Pi, (one probe per chip) are hybridized to the array. Depicted is a probe that has a complementary target on T that is adjacent to the Fx but not to the Fy. Step 3: By applying discriminative conditions or reagents, complexes with no adjacent probes are selectively melted. A particular example is the ligation of a labelled probe to a fixed probe, when the labelled probe hybridizes "back to back" with the attached probe. Positive signals are detected only in the case of adjacent probes, like Fx and Pi, and in a particular example, only in the case of ligated probes.

- 19 -

FIG. 2A, FIG. 2B and FIG. 2C represent components of an exemplary sequencing kit.

FIG. 2A. Sequencing chips, representing an array of  $4^P$  identical sections each containing identical (or  
5 different) arrays of oligonucleotides. Sections can be separated by physical barriers or by hydrophobic strips. 4,000-16,000 oligochips are contemplated to be in the array.

FIG. 2B is an enlargement of a chip section  
10 containing  $4^F$  spots with each with a particular oligonucleotide probe (4,000-16,000) synthesized or spotted on that area. Spots can be as small as several microns and the size of the section is about 1 mm to about 10 mm.

FIG. 2C represents a set of tubes, or one or more  
15 multiwell plates, with an appropriate number of wells (in this case  $4^P$  wells). Each well contains an amount of a specific labeled oligonucleotide. Additional amounts of the probes can be stored unlabeled if the labeling is not  
20 done during synthesis; in this case a sequencing kit will contain necessary components for probe labeling. The lines that are connecting tubes/wells with chip sections depict a step in the sequencing procedure where an amount of a labeled probe is transferred to a chip section. The  
25 transferring can be done by pipetting (single or multi-channel) or by pin array transferring liquid by surface tension. Transferring tools can be also included in the sequencing kit.

FIG. 3A, FIG. 3B and FIG. 3C. Hybridization of DNA  
30 fragments produced by a random cutting of an amount of a DNA molecule. In FIG. 3A, DNA fragment T1 is such that if contains complete targets for both fixed and non-fixed-labeled probes. FIG. 3B represents the case where  
35 the DNA fragment T is not appropriately cut. In FIG. 3C, there is enough space for probe P to hybridize, but the adjacent sequence is not complementary to it. In both

- 20 -

case B and case C, the signal will be reduced due to saturation of the molecules of attached probe F. Simultaneous hybridization with DNA fragments and labeled probes and cycling of the hybridization process are some possible ways to increase the yield of correct adjacent hybridizations.

FIG. 4. Use of Universal Base as a linker or in the terminal position for hybridization. The universal bases (M base, Nichols et al., 1994) or all four bases may be added in the probe synthesis. This is a way to increase the length of the probes, and thus stability of the duplexes without increasing the number of probes. Also the use of universal bases at the free end of probes provides a spacer that allow the sequence to be read in a different frame.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Determining the sequences of nucleic acid molecules is of vital use in all areas of basic and applied biological research (Drmanac & Crkvenjakov, 1990). The present invention provides new and efficient methods for use in sequencing and analyzing nucleic acid molecules. One intended use for this methodology is, in conjunction with other sequencing techniques, for work on the Human Genome Project (HGP).

Presently, two methods of sequencing by hybridization, SBH, are known. In the first, Format 1, unknown genomic DNAs or oligonucleotides of up to about 100-2000 nucleotides in length are arrayed on a solid substrate. These DNAs are then interrogated by hybridization with a set of labeled probes which are generally 6- to 8-mers. In the inverse technique, Format 2, oligomers of 6 to 8 nucleotides are immobilized on a

- 21 -

solid support and allowed to anneal to pieces of cloned and labeled DNA.

5 In either type of SBH analysis, many steps must be included in order to arrive at a definitive sequence. Particular problems of current SBH methods are those associated with the synthesis of large numbers of probes and the difficulties of effective discriminative hybridization. Full match-mismatch discrimination is  
10 difficult due to two main reasons. Firstly, the end mismatch of probes longer than 10 bases is very undiscriminative, and secondly, the complex mixture of labeled DNA segments that result when analyzing a long DNA fragment generates a high background.

15 The present invention provides effective discriminative hybridization without large numbers of probes or probes of increased length, and also eliminates many of the labeling and cloning steps which are  
20 particular disadvantages of each of the known SBH methods. The disclosed highly efficient nucleic acid sequencing methods, termed Format 3 sequencing, are based upon hybridization with two sets of small oligonucleotide probes of known sequences, and thus at least double the  
25 length of sequence that can be determined. These methods allow extremely large nucleic acid molecules, including chromosomes, to be sequenced and solve various other SBH problems such as, for example, the attachment or labelling of many nucleic acid fragments. The invention  
30 is extremely powerful as it may also be used to sequence RNA and even unamplified RNA samples.

Subsequent to the present invention, as disclosed in U.S. Serial No. 08/127,402 and in Drmanac (1994), another  
35 variation of SBH was described termed positional SBH (PSBH) (Broude et al., 1994). PSBH is basically a variant of Format 2 SBH (in which oligonucleotides of

- 22 -

known sequences are immobilized and used to hybridize to nucleic acids of unknown sequence that have been previously labelled). In PSBH, the immobilized probes, rather than being simple, single-stranded probes, are  
5 duplexes that contain single stranded 3' overhangs. Biotinylated duplex probes are immobilized on streptavidin-coated magnetic beads, to form a type of immobilized probe, and then mixed with <sup>32</sup>P-labeled target nucleic acids to be sequenced. T4 DNA ligase is then  
10 added to ligate any hybridized target DNA to the shorter end of the duplex probe.

However, despite representing an interesting approach, PSBH (as reported by Broude et al., 1994) does  
15 not reflect a significant advance over the existing SBH technology. For example, unlike the Format 3 methodology of the present invention, PSBH does not extend the length of sequence that can be determined in one round of the method. PSBH also maintains the burdensome requirement  
20 for labelling the unknown target DNA, which is not required for Format 3. In general, PSBH is proposed for use in comparative studies or in mapping, rather than in de novo genome sequencing. It thus differs significantly from Format 3 which, although widely applicable to all  
25 areas of sequencing, is a very powerful tool for use in sequencing even the largest of genomes.

The nucleic acids to be sequenced may first be fragmented. This may be achieved by any means including,  
30 for example, cutting by restriction enzyme digestion, particularly with Cvi JI as described by Fitzgerald et al. (1992); shearing by physical means such as ultrasound treatment; by NaOH treatment, and the like. If desired, fragments of an appropriate length, such as between about  
35 10 bp and about 40 bp may be cut out of a gel. The complete nucleic acid sequence of the original molecule, such as a human chromosome, would be determined by



- 23 -

defining F + P sequences present in the original molecule and assembling portions of overlapping F + P sequences. This does not, therefore, require an intermediate step of determining fragment sequences, rather, the sequence of the whole molecule is constructed from F + P sequences delineated.

For the purposes of the following discussion, it will be generally assumed that four bases make up the sequences of the nucleic acids to be sequenced. These are A, G, C and T for DNA and A, G, C and U for RNA. However, it may be advantageous in certain embodiments to use modified bases in the small oligonucleotide probes. To carry out the invention, one would generally first prepare a number of small oligonucleotide probes of defined length that cover all combinations of sequences for that length of probe. This number is represented by  $4^N$  (4 to the power N) where the length of the probe is termed N. For example, there are 4096 possible sequences for a 6-mer probe ( $4^6=4096$ ).

One set of such probes of length F ( $4^F$ ) would be fixed in a square array on a microchip - which may be in the range of  $1 \text{ mm}^2$  or  $1 \text{ cm}^2$ . In the present example, these would be arranged in 64 rows and 64 columns. Naturally, one would ensure that the oligo probes were attached, or otherwise immobilized, to the microchip surface so that were able to take part in hybridization reactions. Another set of oligos of length P,  $4^P$  in number, would be also synthesized. The oligos in this "P set" would be labeled with a detectable label and would be dispensed into a set of tubes (FIG. 2A, FIG. 2B and FIG. 2C).

$4^P$  of the chips would be combined in a large array (or several arrays of approximately  $10\text{-}100 \text{ cm}^2$ , for a convenient size); where P corresponds to the length of

- 24 -

oligonucleotides in the second oligomer set (FIG. 2A, FIG. 2B and FIG. 2C). Again, as a convenient example, P is chosen to be six ( $P = 6$ ).

5       The nucleic acids to be sequenced would be fragmented to give smaller nucleic acid fragments of unknown sequence. The average length of these fragments, termed T, should generally be greater than the combined length of F and P and may be about three times the length  
10 of F (i.e.,  $F + P \leq T$  and  $T \approx 3F$ ). In the present example, one would aim to produce nucleic acid fragments of approximately 20 base pairs in length. These fragments would be denatured and added to the large arrays under conditions that facilitate hybridization of  
15 complementary sequences.

In the simplest and currently preferred form of the invention, hybridization conditions would be chosen that would allow significant hybridization to occur only if 6  
20 sequential nucleotides in a nucleic acid fragment were complementary to all 6 nucleotides of an F oligonucleotide probe. Such hybridization conditions would be determined by routine optimization pilot studies in which conditions such as the temperature, the concentration of  
25 various components, the length of time of the steps, and the buffers used, including the pH of the buffer.

At this stage, each microchip would contain certain hybridized complexes. These would be in the form of  
30 probe:fragment complexes in which the entire sequence of the probe is hybridized to the fragment, but in which the fragment, being longer, has some non-hybridized sequences that form a "tail" or "tails" to the complex. In this example, the complementary hybridized sequences would be  
35 of length F and the non-hybridized sequences would total  $T - F$  in length. The complementary portion of the fragment may be at or towards an appropriate end, so that

- 25 -

a single longer non-hybridized tail is formed. Alternatively, the complementary portion of the fragment may be towards the opposite end, so that two non-hybridized tails are formed (FIG. 3A, FIG. 3B and FIG. 3C).

After washing to remove the non-complementary nucleic acid fragments that did not hybridize, a small amount of the labeled oligonucleotides in set P would be added to each microchip for hybridization to the nucleic acid fragment tails of unknown sequence that protrude from the probe:fragment complexes. Only one of each of the 4<sup>P</sup> nucleotides would be added to each microchip. Again, it is currently preferred to use hybridization conditions that would allow significant binding to occur only if all the 6 nucleotides of a labelled probe were complementary to 6 sequential nucleotides of a nucleic acid fragment tail. The hybridization conditions would be determined by pilot studies, as described above, in which components such as the temperature, concentration, time, buffers and the like, are optimized.

At this stage, each microchip would then contain certain 'secondary hybridized complexes'. These would be in the form of probe:fragment:probe complexes in which the entire sequence of each probe is hybridized to the fragment, and in which the fragment likely has some non-hybridized sequences. In these secondary hybridized complexes the immobilized probe and the labelled probe may be hybridized to the fragment so that the two probes are immediately adjacent or "back to back". However, given that the fragments will generally be longer than the sum of the lengths of the probes, the immobilized probe and the labelled probe may be hybridized to the fragment in non-adjacent positions separated by one or more bases.

- 26 -

The large arrays would then be treated by a process to remove the non-hybridized labelled probes. In preferred embodiments, the process employed would remove not only the non-hybridized labelled probes, but also the non-adjacently-hybridized labelled probes from the array. The process would employ discriminating conditions to allow those secondary hybridization complexes that include adjacent immobilized and labelled probes to be discriminating from those secondary hybridization complexes in which the nucleic acid fragment is hybridized to two probes but which probes are not adjacent. This is an important aspect of the invention in that it will allow the ultimate delineation of a section of fragment sequence corresponding to the combined sequences of the immobilized probe and the labelled probe.

The discrimination process employed to remove non-hybridized and non-adjacently-hybridized probes from the array whilst leaving the adjacently-hybridized probes attached may again be a controlled washing process. The adjacently-hybridized probes would be unaffected by the chosen conditions by virtue of their increased stability due to the stacking reactions of the adjacent nucleotides. However, in preferred embodiments, it is contemplated that one would treat the large arrays so that any adjacent probes would be covalently joined, e.g., by treating with a solution containing a chemical ligating agent or, more preferably, a ligase enzyme, such as  $T_4$  DNA ligase (Landegren et al. 1988; Wu & Wallace, 1989).

In any event, the complete array would be subjected to stringent washing so that the only label left associated with the array would be in the form of double-stranded probe-fragment-probe complexes with adjacent hybridized portions of length  $F + P$  (i.e., 12 nucleotides

- 27 -

in the present example). Using this two step hybridization reaction, very high discrimination is possible because three or four independent discriminative processes are taken into account: discriminative hybridization of fragment T to F bases long probe; discriminative hybridization of P bases long probe to fragment T; discriminative stability of full match (F + T + P) hybrid in comparison to P hybrids or even to mismatched hybrids containing non-adjacent F + P probes; and discriminative ligation of the two end bases of F and P.

One would then detect the so-called adjacent secondary complexes by observing the location of the remaining label on the array. From the position of the label, F + P (e.g., 12) nucleotide long sequences from the fragment could be determined by combining the known sequences of the immobilized and labelled probes. The complete nucleic acid sequence of the original molecule, such as a human chromosome, could then be reconstructed or assembled from the overlapping F + P sequences thus determined.

When ligation is employed in the sequencing process, as is currently preferred, then the ordinary oligonucleotides chip cannot be reused. The inventor contemplates that this will not be limiting as various methods are available for recycling. For example, one may generate a specifically cleavable bond between the probes and then cleave the bond after detection. Alternatively, one may employ ribonucleotides for the second probe, probe P, or use a ribonucleotide for the joining base in probe P, so that this probe may subsequently be removed by RNAase or uracil-DNA glycosylate treatment (Craig et al., 1989). Other contemplated methods are to establish bonds by chemical

- 28 -

ligation which can be selectively cut (Dolinnaya et al., 1988).

Further variations and improvements to this sequencing methodology are also contemplated and fall within the scope of the present invention. This includes the use of modified oligonucleotides to increase the specificity or efficiency of the methods, similar to that described by Hoheisel & Lehrach (1990). Cycling hybridizations can also be employed to increase the hybridization signal, as is used in PCR technology. In these cases, one would use cycles with different temperatures to re-hybridize certain probes. The invention also provides for determining shifts in reading frames by using equimolar amounts of probes that have a different base at the end position. For example, using equimolar 7-mers in which the first six bases are the same defined sequence and the last position may be A, T, C or G in the alternative.

20

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples that follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

25

30

35

- 29 -

## EXAMPLE I

PREPARATION OF SUPPORT BOUND OLIGONUCLEOTIDES

5 Oligonucleotides, i.e., small nucleic acid segments, may be readily prepared by, for example, directly synthesizing the oligonucleotide by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer.

10 Support bound oligonucleotides may be prepared by any of the methods known to those of skill in the art using any suitable support such as glass, polystyrene or teflon. One strategy is to precisely spot oligonucleotides synthesized by standard synthesizers.  
15 Immobilization can be achieved using passive adsorption (Inouye & Hondo, 1990); using UV light (Nagata et al., 1985; Dahlen et al., 1987; Morriey & Collins, 1989) or by covalent binding of base modified DNA (Keller et al., 1988; 1989); all references being specifically  
20 incorporated herein.

Another strategy that may be employed is the use of the strong biotin-streptavidin interaction as a linker. For example, Broude et al. (1994) describe the use of  
25 biotinylated probes, although these are duplex probes, that are immobilized on streptavidin-coated magnetic beads. Streptavidin-coated beads may be purchased from Dynal, Oslo. Of course, this same linking chemistry is applicable to coating any surface with streptavidin.  
30 Biotinylated probes may be purchased from various sources, such as, e.g., Operon Technologies (Alameda, CA).

Nunc Laboratories (Naperville, IL) is also selling  
35 suitable material that could be used. Nunc Laboratories have developed a method by which DNA can be covalently bound to the microwell surface termed CovaLink NH.

- 30 -

CovaLink NH is a polystyrene surface grafted with secondary amino groups (>NH) that serve as bridge-heads for further covalent coupling. CovaLink Modules may be purchased from Nunc Laboratories. DNA molecules may be bound to CovaLink exclusively at the 5'-end by a phosphoramidate bond, allowing immobilization of more than 1 pmol of DNA (Rasmussen et al., 1991).

The use of CovaLink NH strips for covalent binding of DNA molecules at the 5'-end has been described (Rasmussen et al., 1991). In this technology, a phosphoramidate bond is employed (Chu et al., 1983). This is beneficial as immobilization using only a single covalent bond is preferred. The phosphoramidate bond joins the DNA to the CovaLink NH secondary amino groups that are positioned at the end of spacer arms covalently grafted onto the polystyrene surface through a 2 nm long spacer arm. To link an oligonucleotide to CovaLink NH via an phosphoramidate bond, the oligonucleotide terminus must have a 5'-end phosphate group. It is, perhaps, even possible for biotin to be covalently bound to CovaLink and then streptavidin used to bind the probes.

More specifically, the linkage method includes dissolving DNA in water (7.5 ng/ $\mu$ l) and denaturing for 10 min. at 95°C and cooling on ice for 10 min. Ice-cold 0.1 M 1-methylimidazole, pH 7.0 (1-MeIm<sub>7</sub>), is then added to a final concentration of 10 mM 1-MeIm<sub>7</sub>. A ss DNA solution is then dispensed into CovaLink NH strips (75  $\mu$ l/well) standing on ice.

Carbodiimide 0.2 M 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC), dissolved in 10 mM 1-MeIm<sub>7</sub>, is made fresh and 25  $\mu$ l added per well. The strips are incubated for 5 hours at 50°C. After incubation the strips are washed using, e.g., Nunc-Immuno Wash; first the wells are washed 3 times, then they are



- 31 -

soaked with washing solution for 5 min., and finally they are washed 3 times (wherein the washing solution is 0.4 N NaOH, 0.25% SDS heated to 50°C).

5           It is contemplated that a further suitable method for use with the present invention is that described in PCT Patent Application WO 90/03382 (Southern & Maskos), incorporated herein by reference. This method of preparing an oligonucleotide bound to a support involves  
10   attaching a nucleoside 3'-reagent through the phosphate group by a covalent phosphodiester link to aliphatic hydroxyl groups carried by the support. The oligonucleotide is then synthesized on the supported nucleoside and protecting groups removed from the  
15   synthetic oligonucleotide chain under standard conditions that do not cleave the oligonucleotide from the support. Suitable reagents include nucleoside phosphoramidite and nucleoside hydrogen phosphorate.

20           In more detail, to use this method, a support, such as a glass plate, is derivatized by contact with a mixture of xylene, glycidoxypropyltrimethoxysilane, and a trace of diisopropylethylamine at 90°C overnight. It is then washed thoroughly with methanol, ether and air-  
25   dried. The derivatized support is then heated with stirring in hexaethyleneglycol containing a catalytic amount of concentrated sulfuric acid, overnight in an atmosphere of argon, at 80°C, to yield an alkyl hydroxyl derivatized support. After washing with methanol and  
30   ether, the support is dried under vacuum and stored under argon at -20°C.

          Oligonucleotide synthesis is then performed by hand under standard conditions using the derivatized glass  
35   plate as a solid support. The first nucleotide will be a 3' - hydrogen phosphate, used in the form of the

- 32 -

triethylammonium salt. This method results in support bound oligonucleotides of high purity.

5 An on-chip strategy for the preparation of DNA probe arrays may be employed. For example, addressable laser-activated photodeprotection may be employed in the chemical synthesis of oligonucleotides directly on a glass surface, as described by Fodor et al. (1991), incorporated herein by reference. Probes may also be  
10 immobilized on nylon supports as described by Van Ness et al. (1991); or linked to teflon using the method of Duncan & Cavalier (1988); all references being specifically incorporated herein.

15 Fodor et al. (1991) describe the light-directed synthesis of dinucleotides which is applicable to the spatially directed synthesis of complex compounds for use in the microfabrication of devices. This is based upon a method that uses light to direct the simultaneous  
20 synthesis of chemical compounds on a solid support. The pattern of exposure to light or other forms of energy through a mask, or by other spatially addressable means, determines which regions of the support are activated for chemical coupling. Activation by light results from the  
25 removal of photolabile protecting groups from selected areas. After deprotection, a first compound bearing a photolabile protecting group is exposed to the entire surface, but reaction occurs only with regions that were addressed by light in the preceding step. The substrate  
30 is then illuminated through a second mask, which activates a different region for reaction with a second protected building block. The pattern of masks used in these illuminations and the sequence of reactants define the ultimate products and their locations. A high degree  
35 of miniaturization is possible with the Fodor method because the density of synthesis sites is bounded only by physical limitations on spatial addressability, i.e., the

- 33 -

diffraction of light. Each compound is accessible and its position is precisely known. hence, an oligo chip made in this way would be ready for use in SBH.

5 Fodor et al. (1991) describes the light-activated formation of a dinucleotide as follows. 5'-Nitroveratryl thymidine was synthesized from the 3'-O-thymidine acetate. After deprotection with base, the 5'-nitroveratryl thymidine was attached to an aminated  
10 substrate through a linkage to the 3'-hydroxyl group. The nitroveratryl protecting groups were removed by illumination through a 500- $\mu$ m checkerboard mask. The substrate was then treated with phosphoramidite-activated 2'-deoxycytidine. In order to follow the reaction  
15 fluorometrically, the deoxycytidine had been modified with an FMOC-protected aminohexyl linker attached to the exocyclic amine. After removal of the FMOC protecting group with base, the regions that contained the dinucleotide were fluorescently labeled by treatment of  
20 the substrate with FITC. Therefore, following this method, support bound-oligonucleotides can be synthesized.

To link an oligonucleotide to a nylon support, as  
25 described by Van Ness et al. (1991), requires activation of the nylon surface via alkylation and selective activation of the 5'-amine of oligonucleotides with cyanuric chloride, as follows. A nylon surface is ethylated using triethyloxonium tetrafluoroborate to form  
30 amine reactive imidate esters on the surface of the nylon 1-methyl-2-pyrrolidone is used as a solvent. The nylon surface is unpolished to effect the greatest possible surface area.

35 The activated surface is then reacted with poly(ethyleneimine) ( $M_r \sim 10K-70K$ ) to form a polymer coating that provides an extended amine surface for the

- 34 -

attachment of oligos. Amine-tailed oligonucleotide(s) selectively react with excess cyanuric chloride, exclusively on the amine tail, to give a 4,6-dichloro-1,3,5-triazinyl-oligonucleotide(s) in quantitative yield.

5 The displacement of one chlorine moiety of cyanuric chloride by the amino group significantly diminishes the reactivity of the remaining chlorine groups. This results in increased hydrolytic stability of the 4,6-dichloro-1,3,5-triazinyl-oligonucleotide(s) are stable

10 for extended periods in buffered aqueous solutions (pH 8.3, 4°C, 1 week) and are readily isolated and purified by size elusion chromatography or ultrafiltration.

The reaction is specific for the amine tail with no

15 apparent reaction on the nucleotide moieties. The PEI-coated nylon surface is then reacted with the cyanuric chloride activated oligonucleotide. High concentrations of the 'capture' sequence are readily immobilized on the surface and the unreacted amines are capped with succinic

20 anhydride in the final step of the derivatization process.

One particular way to prepare support bound oligonucleotides is to utilize the light-generated

25 synthesis described by Pease et al. (1994, incorporated herein by reference). These authors used current photolithographic techniques to generate arrays of immobilized oligonucleotide probes (DNA chips). These methods, in which light is used to direct the synthesis

30 of oligonucleotide probes in high-density, miniaturized arrays, utilize photolabile 5'-protected *N*-acyl-deoxynucleoside phosphoramidites, surface linker chemistry and versatile combinatorial synthesis strategies. A matrix of 256 spatially defined

35 oligonucleotide probes may be generated in this manner and then used in the advantageous Format 3 sequencing, as described herein.

- 35 -

Pease et al. (1994) presented a strategy suitable for use in light-directed oligonucleotide synthesis. In this method, the surface of a solid support modified with photolabile protecting groups is illuminated through a photolithographic mask, yielding reactive hydroxyl groups in the illuminated regions. A 3'-O-phosphoramidite-activated deoxynucleoside (protected at the 5'-hydroxyl with a photolabile group) is then presented to the surface and coupling occurs at sites that were exposed to light. Following capping, and oxidation, the substrate is rinsed and the surface is illuminated through a second mask, to expose additional hydroxyl groups for coupling. A second 5'-protected, 3'-O-phosphoramidite-activated deoxynucleoside is presented to the surface. The selective photodeprotection and coupling cycles are repeated until the desired set of products is obtained. Since photolithography is used, the process can be miniaturized to generate high-density arrays of oligonucleotide probes, the sequence of which is known at each site.

The synthetic pathway for preparing the necessary 5'-O-( $\alpha$ -methyl-6-nitropiperonyloxycabonyl)-N-acyl-2'-deoxynucleoside phosphoramidites (MeNPoc-N-acyl-2'-deoxynucleoside phosphoramidites) involves, in the first step, an N-acyl-2'-deoxynucleoside that reacts with 1-(2-nitro-4,5-methylenedioxyphenyl)ethyl-1-chloroformate to yield 5'-MeNPoc-N-acyl-2'-deoxynucleoside. In the second step, the 3'-hydroxyl reacts with 2-cyanoethyl N,N'-diisopropylchlorophosphoramidite, using standard procedures, to yield the 5'-MeNPoc-N-acyl-2'-deoxynucleoside-3'-O-(2-cyanoethyl-N,N'-diisopropyl)phosphoramidites. The photoprotecting group is stable under ordinary phosphoramidite synthesis conditions and can be removed with aqueous base. These reagents can be stored for long periods under argon at 4°C.

- 36 -

Photolysis half-times of 28 s, 31 s, 27 s, and 18 s for MeNPoc-dT, MeNPoc-dC<sup>ibu</sup>, MeNPoc-dG<sup>PAC</sup>, and MeNPoc-dA<sup>PAC</sup> respectively, have been reported (Pease et al., 1994). In lithographic synthesis, illumination times of 4.5 min (9 x t<sub>1/2</sub> MeNPoc-dC) are therefore recommended to ensure >99% removal of MeNPoc protecting groups.

A suitable synthetic support is one consisting of a 5.1 x 7.6 cm glass substrate prepared by cleaning in concentrated NaOH, followed by exhaustive rinsing in water. The surfaces would then be derivatized for 2 hr with a solution of 10% (vol/vol) bis(2-hydroxyethyl)aminopropyltriethoxysilane (Petrarch Chemicals, Bristol, PA) in 95% ethanol, rinsed thoroughly with ethanol and ether, dried in vacuo at 40°C, and heated at 100°C for 15 min. In such studies, a synthesis linker would be attached by reacting derivatized substrates with 4,4'-dimethoxytrityl (DMT)-hexaethyloxy-O-cyanoethyl phosphoramidite.

In summary, to initiate the synthesis of an oligonucleotide probe, the appropriate deoxynucleoside phosphoramidite derivative would be attached to a synthetic support through a linker. Regions of the support are then activated for synthesis by illumination through, e.g., 800 x 12800 µm apertures of a photolithographic mask. Additional phosphoramidite synthesis cycles may be performed (with DMT-protected deoxynucleosides) to generate any required sequence, such as any 4-, 5-, 6-, 7-, 8-, 9- or even 10-mer sequence. Following removal of the phosphate and exocyclic amine protecting groups with concentrated NH<sub>4</sub>OH for 4 hr, the substrate may then be mounted in a water-jacketed thermostatically controlled hybridization chamber, ready for use.

- 37 -

Of course, one could easily purchase a DNA chip, such as one of the light-activated chips described above, from a commercial source. In this regard, one may contact Affymetrix of Santa Clara, CA 95051, and  
5 Beckman.

## EXAMPLE II

### MODIFIED OLIGONUCLEOTIDES FOR USE IN PROBES

10

Modified oligonucleotides may be used throughout the procedures of the present invention to increase the specificity or efficiency of hybridization. A way to achieve this is the substitution of natural nucleotides  
15 by base modification. For example, pyrimidines with a halogen at the C<sup>5</sup>-position may be used. This is believed to improve duplex stability by influencing base stacking. 2,6-diaminopurine may also be used to give a third  
20 hydrogen bond in its base pairing with thymine, thereby thermally stabilizing DNA-duplexes. Using 2,6-diaminopurine is reported to lead to a considerable improvement in the duplex stability of short oligomers. Its incorporation is proposed to allow more stringent conditions for primer annealing, thereby improving the  
25 specificity of the duplex formation and suppressing background problems or the use of shorter oligomers.

The synthesis of the triphosphate versions of these modified nucleotides is disclosed by Hoheisel & Lehrach  
30 (1990, incorporated herein by reference). Briefly, 5-Chloro-2'-deoxyuridine and 2,6-diaminopurine 2'-deoxynucleoside are purchased, e.g., from Sigma. Phosphorylation is carried out as follows: 50 mg dry 2-NH<sub>2</sub>-dAdo is taken up in 500  $\mu$ l dry triethyl phosphate  
35 stirring under argon. 25  $\mu$ l POCl<sub>3</sub> is added and the mixture incubated at -20°C. In the meantime, 1 mmol pyrophosphoric acid is dissolved in 0.95 ml tri-n-

- 38 -

butylamine and 2 ml methanol and dried in a rotary evaporator. Subsequently it is dried by evaporation twice from 5 ml pyridine, with 70  $\mu$ l tri-n-butylamine also added before the second time. Finally it is  
5 dissolved in 2 ml dry dimethyl formamide.

After 90 min at  $-20^{\circ}\text{C}$ , the phosphorylation mixture is evaporated to remove excess  $\text{POCl}_3$  and the tri-n-butylammonium pyrophosphate in dimethyl formamide is  
10 added. Incubation is for 1.5 min at room temperature. The reaction is stopped by addition of 5 ml 0.2 M triethylammonium bicarbonate (pH 7.6) and kept on ice for 4 hours. For 5-Cl-dUrd, the conditions would be identical, but 50  $\mu$ l  $\text{POCl}_3$  would be added and the  
15 phosphorylation carried out at room temperature for 4 hours.

After the hydrolysis, the mixture is evaporated, the pH adjusted to 7.5, and extracted with 1 volume diethyl  
20 ether. Separation of the products is, e.g., on a (2.5 x 20 cm) Q-Sepharose column using a linear gradient of 0.15 M to 0.8 M triethylammonium bicarbonate. Stored frozen, the nucleotides are stable over long periods of time.

25 One may also use the non-discriminatory base analogue, or universal base, as designed by Nichols et al. (1994). This new analogue, 1-(2'-deoxy- $\beta$ -D-ribofuranosyl)-3-nitropyrrole (designated M), was generated for use in oligonucleotide probes and primers  
30 for solving the design problems that arise as a result of the degeneracy of the genetic code, or when only fragmentary peptide sequence data are available. This analogue maximizes stacking while minimizing hydrogen-bonding interactions without sterically disrupting a DNA  
35 duplex.



- 39 -

The M nucleoside analogue was designed to maximize stacking interactions using aprotic polar substituents linked to heteroaromatic rings, enhancing intra- and inter-strand stacking interactions to lessen the role of hydrogen bonding in base-pairing specificity. Nichols et al. (1994) favored 3-nitropyrrole 2'-deoxyribonucleoside because of its structural and electronic resemblance to *p*-nitroaniline, whose derivatives are among the smallest known intercalators of double-stranded DNA.

The dimethoxytrityl-protected phosphoramidite of nucleoside M is also available for incorporation into nucleotides used as primers for sequencing and polymerase chain reaction (PCR). Nichols et al. (1994) showed that a substantial number of nucleotides can be replaced by M without loss of primer specificity.

A unique property of M is its ability to replace long strings of contiguous nucleosides and still yield functional sequencing primers. Sequences with three, six and nine M substitutions have all been reported to give readable sequencing ladders, and PCR with three different M-containing primers all resulted in amplification of the correct product (Nichols et al., 1994).

The ability of 3-nitropyrrole-containing oligonucleotides to function as primers strongly suggests that a duplex structure must form with complementary strands. Optical thermal profiles obtained for the oligonucleotide pairs d(5'-C<sub>2</sub>-T<sub>5</sub>XT<sub>5</sub>G<sub>2</sub>-3') and d(5'-C<sub>2</sub>A<sub>5</sub>YA<sub>5</sub>G<sub>2</sub>-3') (where X and Y can be A, C, G, T or M) were reported to fit the normal sigmoidal pattern observed for the DNA double-to-single strand transition. The *T<sub>m</sub>* values of the oligonucleotides containing X·M base pairs (where X was A, C, G or T, and Y was M) were reported to all fall within a 3°C range (Nichols et al., 1994).

- 40 -

## EXAMPLE III

PREPARATION OF SEQUENCING CHIPS AND ARRAYS

5           The present example describes physical embodiments of sequencing chips contemplated by the inventor.

          A basic example is using 6-mers attached to 50 micron surfaces to give a chip with dimensions of  
10   3 x 3 mm which can be combined to give an array of 20 x 20 cm. Another example is using 9-mer oligonucleotides attached to 10 x 10 microns surface to create a 9-mer chip, with dimensions of 5 x 5 mm. 4000 units of such chips may be used to create a 30 x 30 cm  
15   array. FIG. 2A, FIG. 2B and FIG. 2C illustrate yet another example of an array in which 4,000 to 16,000 oligochips are arranged into a square array. A plate, or collection of tubes, as also depicted, may be packaged with the array as part of the sequencing kit.

20

          The arrays may be separated physically from each other or by hydrophobic surfaces. One possible way to utilize the hydrophobic strip separation is to use technology such as the Iso-Grid Microbiology System  
25   produced by QA Laboratories, Toronto, Canada.

          Hydrophobic grid membrane filters (HGMF) have been in use in analytical food microbiology for about a decade where they exhibit unique attractions of extended  
30   numerical range and automated counting of colonies. One commercially-available grid is ISO-GRID™ from QA Laboratories Ltd. (Toronto, Canada) which consists of a square (60 x 60 cm) of polysulfone polymer (Gelman Tuffryn HT-450, 0.45μ pore size) on which is printed a  
35   black hydrophobic ink grid consisting of 1600 (40 x 40) square cells. HGMF have previously been inoculated with

- 41 -

bacterial suspensions by vacuum filtration and incubated on the differential or selective media of choice.

Because the microbial growth is confined to grid cells of known position and size on the membrane, the HGMF functions more like an MPN apparatus than a conventional plate or membrane filter. Peterkin et al. (1987) reported that these HGMFs can be used to propagate and store genomic libraries when used with a HGMF replicator. One such instrument replicates growth from each of the 1600 cells of the ISO-GRID and enables many copies of the master HGMF to be made (Peterkin et al., 1987).

Sharpe et al. (1989) also used ISO-GRID HGMF from QA Laboratories and an automated HGMF counter (MI-100 Interpreter) and RP-100 Replicator. They reported a technique for maintaining and screening many microbial cultures.

Peterkin and colleagues later described a method for screening DNA probes using the hydrophobic grid-membrane filter (Peterkin et al., 1989). These authors reported methods for effective colony hybridization directly on HGMFs. Previously, poor results had been obtained due to the low DNA binding capacity of the polysulfone polymer on which the HGMFs are printed. However, Peterkin et al. (1989) reported that the binding of DNA to the surface of the membrane was improved by treating the replicated and incubated HGMF with polyethyleneimine, a polycation, prior to contact with DNA. Although this early work uses cellular DNA attachment, and has a different objective to the present invention, the methodology described may be readily adapted for format 3 SBH.

35

In order to identify useful sequences rapidly, Peterkin et al. (1989) used radiolabeled plasmid DNA from

- 42 -

various clones and tested its specificity against the DNA on the prepared HGMFs. In this way, DNA from recombinant plasmids was rapidly screened by colony hybridization against 100 organisms on HGMF replicates which can be easily and reproducibly prepared.

Two basic problems have to be solved. Manipulation with small (2-3 mm) chips, and parallel execution of thousands of the reactions. The solution of the invention is to keep the chips and the probes in the corresponding arrays. In one example, chips containing 250,000 9-mers are synthesized on a silicon wafer in the form of 8x8 mm plates (15  $\mu$ M/oligonucleotide, Pease et al., 1994) arrayed in 8x12 format (96 chips) with a 1 mm groove in between. Probes are added either by multichannel pipet or pin array, one probe on one chip. To score all 4000 6-mers, 42 chip arrays have to be used, either using different ones, or by reusing one set of chip arrays several times.

20

In the above case, using the earlier nomenclature of the application,  $F = 9$ ;  $P = 6$ ; and  $F + P = 15$ . Chips may have probes of formula  $B_xN_n$ , where  $x$  is a number of specified bases  $B$ ; and  $n$  is a number of non-specified bases, so that  $x = 4$  to 10 and  $n = 1$  to 4. To achieve more efficient hybridization, and to avoid potential influence of any support oligonucleotides, the specified bases can be surrounded by unspecified bases, thus represented by a formula such as  $(N)_nB_x(N)_m$  (FIG. 4).

30

#### EXAMPLE IV

##### PREPARATION OF NUCLEIC ACID FRAGMENTS

The nucleic acids to be sequenced may be obtained from any appropriate source, such as cDNAs, genomic DNA, chromosomal DNA, microdissected chromosome bands, cosmid

- 43 -

or YAC inserts, and RNA, including mRNA without any amplification steps. For example, Sambrook et al. (1989) describes three protocols for the isolation of high molecular weight DNA from mammalian cells (p. 9.14-9.23).

5

The nucleic acids would then be fragmented by any of the methods known to those of skill in the art including, for example, using restriction enzymes as described at 9.24-9.28 of Sambrook et al. (1989), shearing by ultrasound and NaOH treatment.

Low pressure shearing is also appropriate, as described by Schrieffer et al. (1990, incorporated herein by reference). In this method, DNA samples are passed through a small French pressure cell at a variety of low to intermediate pressures. A lever device allows controlled application of low to intermediate pressures to the cell. The results of these studies indicate that low-pressure shearing is a useful alternative to sonic and enzymatic DNA fragmentation methods.

One particularly suitable way for fragmenting DNA is contemplated to be that using the two base recognition endonuclease, CviJI, described by Fitzgerald et al. (1992). These authors described an approach for the rapid fragmentation and fractionation of DNA into particular sizes that they contemplated to be suitable for shotgun cloning and sequencing. The present inventor envisions that this will also be particularly useful for generating random, but relatively small, fragments of DNA for use in the present sequencing technology.

The restriction endonuclease CviJI normally cleaves the recognition sequence PuGCPy between the G and C to leave blunt ends. Atypical reaction conditions, which alter the specificity of this enzyme (CviJI\*\*), yield a

- 44 -

quasi-random distribution of DNA fragments from the small molecule pUC19 (2688 base pairs). Fitzgerald et al. (1992) quantitatively evaluated the randomness of this fragmentation strategy, using a CviJI\*\* digest of pUC19 that was size fractionated by a rapid gel filtration method and directly ligated, without end repair, to a lacZ minus M13 cloning vector. Sequence analysis of 76 clones showed that CviJI\*\* restricts PyGCPy and PuGCPu, in addition to PuGCPy sites, and that new sequence data is accumulated at a rate consistent with random fragmentation.

As reported in the literature, advantages of this approach compared to sonication and agarose gel fractionation include: smaller amounts of DNA are required (0.2-0.5  $\mu$ g instead of 2-5  $\mu$ g); and fewer steps are involved (no preligation, end repair, chemical extraction, or agarose gel electrophoresis and elution are needed). These advantages are also proposed to be of use when preparing DNA for sequencing by Format 3.

Irrespective of the manner in which the nucleic acid fragments are obtained or prepared, it is important to denature the DNA to give single stranded pieces available for hybridization. This is achieved by incubating the DNA solution for 2-5 minutes at 80-90°C. The solution is then cooled quickly to 2°C to prevent renaturation of the DNA fragments before they are contacted with the chip. Phosphate groups must also be removed from genomic DNA, as described in Example VI.

#### EXAMPLE V

##### PREPARATION OF LABELLED PROBES

The oligonucleotide probes may be prepared by automated synthesis, which is routine to those of skill in the art, for example, using an Applied Biosystems system. Alternatively, probes may be prepared using

- 45 -

Genosys Biotechnologies Inc. methods using stacks of porous Teflon wafers.

Oligonucleotide probes may be labelled with, for example, radioactive labels ( $^{35}\text{S}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ , and preferably,  $^{33}\text{P}$ ) for arrays with 100-200  $\mu\text{m}$  or 100-400  $\mu\text{m}$  spots; non-radioactive isotopes (Jacobsen et al., 1990); or fluorophores (Brumbaugh et al., 1988). All such labelling methods are routine in the art, as exemplified by the relevant sections in Sambrook et al. (1989) and by further references such as Schubert et al. (1990), Murakami et al. (1991) and Cate et al. (1991), all articles being specifically incorporated herein by reference.

15

In regard to radiolabeling, the common methods are end-labelling using T4 polynucleotide kinase or high specific activity labelling using Klenow or even T7 polymerase. These are described as follows.

20

Synthetic oligonucleotides are synthesized without a phosphate group at their 5' termini and are therefore easily labeled by transfer of the  $\gamma$ - $^{32}\text{P}$  or  $\gamma$ - $^{33}\text{P}$  from  $[\gamma$ - $^{32}\text{P}]\text{ATP}$  or  $[\gamma$ - $^{33}\text{P}]\text{ATP}$  using the enzyme bacteriophage T4 polynucleotide kinase. If the reaction is carried out efficiently, the specific activity of such probes can be as high as the specific activity of the  $[\gamma$ - $^{32}\text{P}]\text{ATP}$  or  $[\gamma$ - $^{33}\text{P}]\text{ATP}$  itself. The reaction described below is designed to label 10 pmoles of an oligonucleotide to high specific activity. Labeling of different amounts of oligonucleotide can easily be achieved by increasing or decreasing the size of the reaction, keeping the concentrations of all components constant.

30

A reaction mixture would be created using 1.0  $\mu\text{l}$  of oligonucleotide (10 pmoles/ $\mu\text{l}$ ); 2.0  $\mu\text{l}$  of 10  $\times$  bacteriophage T4 polynucleotide kinase buffer; 5.0  $\mu\text{l}$  of

35

- 46 -

[ $\gamma$ - $^{32}\text{P}$ ]ATP or [ $\gamma$ - $^{33}\text{P}$ ]ATP (sp. act. 5000 Ci/mmol; 10 mCi/ml in aqueous solution) (10 pmoles); and 11.4  $\mu\text{l}$  of water. Eight (8) units ( $\sim 1 \mu\text{l}$ ) of bacteriophage T4 polynucleotide kinase is added to the reaction mixture  
5 mixed well, and incubated for 45 minutes at 37°C. The reaction is heated for 10 minutes at 68°C to inactivate the bacteriophage T4 polynucleotide kinase.

The efficiency of transfer of  $^{32}\text{P}$  or  $^{33}\text{P}$  to the  
10 oligonucleotide and its specific activity is then determined. If the specific activity of the probe is acceptable, it is purified. If the specific activity is too low, an additional 8 units of enzyme is added and incubated for a further 30 minutes at 37°C before heating  
15 the reaction for 10 minutes at 68°C to inactivate the enzyme.

Purification of radiolabeled oligonucleotides can be achieved by precipitation with ethanol; precipitation  
20 with cetylpyridinium bromide; by chromatography through bio-gel P-60; or by chromatography on a Sep-Pak C<sub>18</sub> column.

Probes of higher specific activities can be obtained  
25 using the Klenow fragment of *E. coli*. DNA polymerase I to synthesize a strand of DNA complementary to the synthetic oligonucleotide. A short primer is hybridized to an oligonucleotide template whose sequence is the complement of the desired radiolabeled probe. The primer  
30 is then extended using the Klenow fragment of *E. coli* DNA polymerase I to incorporate [ $\alpha$ - $^{32}\text{P}$ ]dNTPs or [ $\alpha$ - $^{33}\text{P}$ ]dNTPs in a template-directed manner. After the reaction, the template and product are separated by denaturation followed by electrophoresis through a polyacrylamide gel  
35 under denaturing conditions. With this method, it is possible to generate oligonucleotide probes that contain



- 47 -

several radioactive atoms per molecule of oligonucleotide, if desired.

To use this method, one would mix in a microfuge  
5 tube the calculated amounts of [ $\alpha$ -<sup>32</sup>P]dNTPs or [ $\alpha$ -<sup>33</sup>P]dNTPs necessary to achieve the desired specific activity and sufficient to allow complete synthesis of all template strands. The concentration of dNTPs should not be less than 1 $\mu$ M at any stage during the reaction.  
10 Then add to the tube the appropriate amounts of primer and template DNAs, with the primer being in three- to tenfold molar excess over the template.

0.1 volume of 10  $\times$  Klenow buffer would then be added  
15 and mixed well. 2-4 units of the Klenow fragment of *E. coli* DNA polymerase I would then be added per 5  $\mu$ l of reaction volume, mixed and incubated for 2-3 hours at 4°C. If desired, the progress of the reaction may be monitored by removing small (0.1- $\mu$ l) aliquots and  
20 measuring the proportion of radioactivity that has become precipitable with 10% trichloroacetic acid (TCA).

The reaction would be diluted with an equal volume of gel-loading buffer, heated to 80°C for 3 minutes, and  
25 then the entire sample loaded on a denaturing polyacrylamide gel. Following electrophoresis, the gel is autoradiographed, allowing the probe to be localized and removed from the gel. Various methods for fluorophobic labelling are also available, as follows.  
30 Brumbaugh et al. (1988) describe the synthesis of fluorescently labeled primers. A deoxyuridine analog with a primary amine "linker arm" of 12 atoms attached at C-5 is synthesized. Synthesis of the analog consists of derivatizing 2'-deoxyuridine through organometallic  
35 intermediates to give 5' (methyl propenoyl)-2'-deoxyuridine. Reaction with dimethoxytrityl-chloride produces the corresponding 5'-dimethoxytrityl adduct.

- 48 -

The methyl ester is hydrolyzed, activated, and reacted with an appropriately monoacylated alkyl diamine. After purification, the resultant linker arm nucleosides are converted to nucleoside analogs suitable for chemical  
5 oligonucleotide synthesis.

Oligonucleotides would then be made that include one or two linker arm bases by using modified phosphoridite chemistry. To a solution of 50 nmol of the linker arm  
10 oligonucleotide in 25  $\mu$ l of 500 mM sodium bicarbonate (pH 9.4) is added 20  $\mu$ l of 300 mM FITC in dimethyl sulfoxide. The mixture is agitated at room temperature for 6 hr. The oligonucleotide is separated from free FITC by elution from a 1 x 30 cm Sephadex G-25 column with 20 mM  
15 ammonium acetate (pH 6), combining fractions in the first UV-absorbing peak.

In general, fluorescent labelling of an oligonucleotide at its 5'-end initially involved two  
20 steps. First, a N-protected aminoalkyl phosphoramidite derivative is added to the 5'-end of an oligonucleotide during automated DNA synthesis. After removal of all protecting groups, the NHS ester of an appropriate fluorescent dye is coupled to the 5'-amino group  
25 overnight followed by purification of the labelled oligonucleotide from the excess of dye using reverse phase HPLC or PAGE.

Schubert et al. (1990) described the synthesis of a  
30 phosphoramidite that enables oligonucleotides labeled with fluorescein to be produced during automated DNA synthesis. Fluorescein methylester is alkylated with 4-chloro(4,4'-dimethoxytrityl)butanol-1 in the presence of  $K_2CO_3$  and KI in DMF for 17 hrs. After removal of the  
35 trityl group with 1% TFA in chloroform, the product is phosphitylated by standard procedures with bis(diisopropylamino)methoxyphosphine. Phosphorylation

- 49 -

of the above obtained fluorescein derivative leads an H-phosphonate in reasonable yields. The resulting amidite (0.1 M solution in dry acetonitrile) is used for the automated synthesis of different primers using  $\beta$ -cyanoethyl phosphoramidite chemistry and a DNA synthesizer. Cleavage from the support and deprotection is performed with 25% aqueous ammonia for 36 hrs at room temperature. The crude product is purified by PAGE and the labelled primer is visible as a pale green fluorescent band at 310 nm. Elution and desalting using RP 18 cartridges yields the desired product.

The fluorescent labelling of the 5'-end of a probe in the Schubert method is directly achieved during DNA synthesis in the last coupling cycle. Coupling yields are as high as with the normal phosphoramidites. After deprotection and removal of ammonia by lyophilization using a speed vac or by ethanol precipitation, fluorescent labelled oligonucleotides can be directly used for DNA sequencing in Format 3 SBH.

Murakami et al. also described the preparation of fluorescein-labeled oligonucleotides. This synthesis is based on a polymer-supported phosphoramidite and hydrogen phosphonate method. Ethylenediamine or hexamethylenediamine is used as a tether. They were introduced via a phosphoramidate linkage, which was formed by oxidation of a hydrogen-phosphonate intermediate in  $\text{CCl}_4$  solution. The modified oligonucleotides are subjected to labeling using a primary amine orienting reagent, FITC, on the beads. The resulting modified oligonucleotide is cleaved from beads and subsequently purified by RPLC.

Cate et al. (1991) describe the use of oligonucleotide probes directly conjugated to alkaline phosphatase in combination with a direct chemiluminescent

- 50 -

substrate (AMPPD) to allow probe detection. Alkaline phosphatase may be covalently coupled to a modified base of the oligonucleotide. After hybridization, the oligo would be incubated with AMPDD. The alkaline phosphatase enzyme breaks AMPDD to yield a compound that produces fluorescence without excitation, i.e., a laser is not needed. It is contemplated that a strong signal can be generated using such technology.

Labelled probes could readily be purchased from a variety of commercial sources, including GEN<sub>5</sub>ET, rather than synthesized.

15

#### EXAMPLE VI

##### REMOVAL OF PHOSPHATE GROUPS

Both bacterial alkaline phosphatase (BAP) and calf intestinal alkaline phosphatase (CIP) catalyze the removal of 5'-phosphate residues from DNA and RNA. They are therefore appropriate for removing 5' phosphates from DNA and/or RNA to prevent ligation and inappropriate hybridization. Phosphate removal, as described by Sambrook et al. (1989), would be performed after cutting, or otherwise shearing, the genomic DNA.

BAP is the more active of the two alkaline phosphatases, but it is also far more resistant to heat and detergents. It is therefore difficult to inhibit BAP completely at the end of dephosphorylation reactions. Proteinase K is used to digest CIP, which must be completely removed if subsequent ligations are to work efficiently. An alternative method is to inactivate the CIP by heating to 65°C for 1 hour (or 75°C for 10 minutes) in the presence of 5 mM EDTA (pH 8.0) and then to purify the dephosphorylated DNA by extraction with phenol:chloroform.

## EXAMPLE VII

CONDUCTING SEQUENCING BY TWO STEP HYBRIDIZATION

5           Following are certain examples to describe the  
execution of the sequencing methodology contemplated by  
the inventor. First, the whole chip would be hybridized  
with mixture of DNA as complex as 100 million of bp (one  
human chromosome). Guidelines for conducting  
10   hybridization can be found in papers such as Drmanac  
et al. (1990); Khrapko et al. (1991); and Broude et al.  
(1994). These articles teach the ranges of hybridization  
temperatures, buffers and washing steps that are  
appropriate for use in the initial step of Format 3 SBH.

15           The present inventor particularly contemplates that  
hybridization is to be carried out for up to several  
hours in high salt concentrations at a low temperature  
(-2°C to 5°C) because of a relatively low concentration  
20   of target DNA that can be provided. For this purpose,  
SSC buffer is used instead of sodium phosphate buffer  
(Drmanac et al., 1990), which precipitates at 10°C.  
Washing does not have to be extensive (a few minutes)  
because of the second step, and can be completely  
25   eliminated when the hybridization cycling is used for the  
sequencing of highly complex DNA samples. The same  
buffer is used for hybridization and washing steps to be  
able to continue with the second hybridization step with  
labeled probes.

30           After proper washing using a simple robotic device  
on each array, e.g., a 8 x 8mm array (Example III), one  
labeled, probe, e.g., a 6-mer, would be added. A 96-tip  
or 96-pin device would be used, performing this in 42  
35   operations. Again, a range of discriminatory conditions  
could be employed, as previously described in the  
scientific literature.

- 52 -

The present inventor particularly contemplates the use of the following conditions. First, after adding labeled probes and incubating for several minutes only (because of the high concentration of added  
5 oligonucleotides) at a low temperature (0-5°C), the temperature is increased to 3-10°C, depending on F+P length, and the washing buffer is added. At this time, the washing buffer used is one compatible with any ligation reaction (e.g., 100 mM salt concentration  
10 range). After adding ligase, the temperature is increased again to 15-37°C to allow fast ligation (less than 30 min) and further discrimination of full match and mismatch hybrids.

15 The use of cationic detergents is also contemplated for use in Format 3 SBH, as described by Pontius & Berg (1991, incorporated herein by reference). These authors describe the use of two simple cationic detergents, dedecyl- and cetyltrimethylammonium bromide (DTAB and  
20 CTAB) in DNA renaturation.

DTAB and CTAB are variants of the quaternary amine tetramethylammonium bromide (TMAB) in which one of the methyl groups is replaced by either a 12-carbon (DTAB) or  
25 a 16-carbon (CTAB) alkyl group. TMAB is the bromide salt of the tetramethylammonium ion, a reagent used in nucleic acid renaturation experiments to decrease the G-C-content bias of the melting temperature. DTAB and CTAB are similar in structure to sodium dodecyl sulfate (SDS),  
30 with the replacement of the negatively charged sulfate of SDS by a positively charged quaternary amine. While SDS is commonly used in hybridization buffers to reduce nonspecific binding and inhibit nucleases, it does not greatly affect the rate of renaturation.

35

- 53 -

When using a ligation process, the enzyme could be added with the labeled probes or after the proper washing step to reduce the background.

5           Although not previously proposed for use in any SBH method, ligase technology is well established within the field of molecular biology. For example, Hood and colleagues described a ligase-mediated gene detection technique (Landegren et al., 1988), the methodology of  
10           which can be readily adapted for use in Format 3 SBH. Landegren et al. describe an assay for the presence of given DNA sequences based on the ability of two oligonucleotides to anneal immediately adjacent to each other on a complementary target DNA molecule. The two  
15           oligonucleotides are then joined covalently by the action of a DNA ligase, provided that the nucleotides at the junction are correctly base-paired. Although not previously contemplated, this situation now arises in Format 3 sequencing. Wu & Wallace also describe the use  
20           of bacteriophage T4 DNA ligase to join two adjacent, short synthetic oligonucleotides. Their oligo ligation reactions were carried out in 50 mM Tris HCl pH 7.6, 10 mM MgCl<sub>2</sub>, 1 mM ATP, 1 mM DTT, and 5% PEG. Ligation reactions were heated to 100°C for 5-10 min followed by  
25           cooling to 0°C prior to the addition of T4 DNA ligase (1 unit; Bethesda Research Laboratory). Most ligation reactions were carried out at 30°C and terminated by heating to 100°C for 5 min.

30           Final washing appropriate for discriminating detection of hybridized adjacent, or ligated, oligonucleotides of length (F + P), is then performed. This washing step is done in water for several minutes at 40-60°C to wash out all the non ligated labeled probes,  
35           and all other compounds, to maximally reduce background. Because of the covalently bound labeled oligonucleotides,

- 54 -

detection is simplified (it does not have time and low temperature constraints).

Depending on the label used, imaging of the chips is done with different apparatus. For radioactive labels, phosphor storage screen technology and PhosphorImager as a scanner may be used (Molecular Dynamics, Sunnyvale, CA). Chips are put in a cassette and covered by a phosphorous screen. After 1-4 hours of exposure, the screen is scanned and the image file stored at a computer hard disc. For the detection of fluorescent labels, CCD cameras and epifluorescent or confocal microscopy are used. For the chips generated directly on the pixels of a CCD camera, detection can be performed as described by Eggers et al. (1994, incorporated herein by reference).

Charge-coupled device (CCD) detectors serve as active solid supports that quantitatively detect and image the distribution of labeled target molecules in probe-based assays. These devices use the inherent characteristics of microelectronics that accommodate highly parallel assays, ultrasensitive detection, high throughput, integrated data acquisition and computation. Eggers et al. (1994) describe CCDs for use with probe-based assays, such as Format 3 SBH of the present invention, that allow quantitative assessment within seconds due to the high sensitivity and direct coupling employed.

The integrated CCD detection approach enables the detection of molecular binding events on chips. The detector rapidly generates a two-dimensional pattern that uniquely characterizes the sample. In the specific operation of the CCD-based molecular detector, distinct biological probes are immobilized directly on the pixels of a CCD or can be attached to a disposable cover slip placed on the CCD surface. The sample molecules can be



- 55 -

labeled with radioisotope, chemiluminescent or fluorescent tags.

Upon exposure of the sample to the CCD-based probe  
5 array, photons or radioisotope decay products are emitted  
at the pixel locations where the sample has bound, in the  
case of Format 3, to two complementary probes. In turn,  
electron-hole pairs are generated in the silicon when the  
10 charged particles, or radiation from the labeled sample,  
are incident on the CCD gates. Electrons are then  
collected beneath adjacent CCD gates and sequentially  
read out on a display module. The number of  
photoelectrons generated at each pixel is directly  
proportional to the number of molecular binding events in  
15 such proximity. Consequently, molecular binding can be  
quantitatively determined (Eggers et al., 1994).

As recently reported, silicon-based CCDs have  
advantages as solid-state detection and imaging sensors  
20 primarily because of the high sensitivity of the devices  
over a wide wavelength range (from 1 to 10000 Å).  
Silicon is very responsive to electromagnetic radiation  
from the visible spectrum to soft X-rays. For visible  
light, a single photon incident on the CCD gate results  
25 in a single electron charge packet beneath the gate. A  
single soft X-ray beta particle (typically KeV to MeV  
range) generates thousands to tens of thousands of  
electrons. In addition to the high sensitivity, the CCDs  
described by Eggers et al. (1994) offer a wide dynamic  
30 range (4 to 5 orders of magnitude) since a detectable  
charge packet can range from a few to  $10^5$  electrons. The  
detection response is linear over a wide dynamic range.

By placing the imaging array in proximity to the  
35 sample, the collection efficiency is improved by a factor  
of at least 10 over lens-based techniques such as those  
found in conventional CCD cameras. That is, the sample

- 56 -

(emitter) is in near contact with the detector (imaging array), and this eliminates conventional imaging optics such as lenses and mirrors.

5           When radioisotopes are attached as reporter groups to the target molecules, energetic particles are detected. Several reporter groups that emit particles of varying energies have been successfully utilized with the micro-fabricated detectors, including  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$   
10   and  $^{125}\text{I}$ . The higher energy particles, such as from  $^{32}\text{P}$ , provide the highest molecular detection sensitivity, whereas the lower energy particles, such as from  $^{35}\text{S}$ , provide better resolution. Hence, the choice of the radioisotope reporter can be tailored as required. Once  
15   the particular radioisotope label is selected, the detection performance can be predicted by calculating the signal-to-noise ratio (SNR), as described by Eggers *et al.* (1994).

20           An alternative luminescent detection procedure involves the use of fluorescent or chemiluminescent reporter groups attached to the target molecules. The fluorescent labels can be attached covalently or through interaction. Fluorescent dyes, such as ethidium bromide,  
25   with intense absorption bands in the near UV (300-350 nm) range and principal emission bands in the visible (500-650 nm) range, are most suited for the CCD devices employed since the quantum efficiency is several orders of magnitude lower at the excitation wavelength than at  
30   the fluorescent signal wavelength.

          From the perspective of detecting luminescence, the polysilicon CCD gates have the built-in capacity to filter away the contribution of incident light in the UV  
35   range, yet are very sensitive to the visible luminescence generated by the fluorescent reporter groups. Such inherently large discrimination against UV excitation

- 57 -

enables large SNRs (greater than 100) to be achieved by the CCDs as formulated in the incorporated paper by Eggers et al. (1994).

5           For probe immobilization on the detector, hybridization matrices may be produced on inexpensive  $\text{SiO}_2$  wafers, which are subsequently placed on the surface of the CCD following hybridization and drying. This format is economically efficient since the hybridization  
10 of the DNA is conducted on inexpensive disposable  $\text{SiO}_2$  wafers, thus allowing reuse of the more expensive CCD detector. Alternatively, the probes can be immobilized directly on the CCD to create a dedicated probe matrix.

15           To immobilize probes upon the  $\text{SiO}_2$  coating, a uniform epoxide layer is linked to the film surface, employing an epoxy-silane reagent and standard  $\text{SiO}_2$  modification chemistry. Amine-modified oligonucleotide probes are then linked to the  $\text{SiO}_2$  surface by means of  
20 secondary amine formation with the epoxide ring. The resulting linkage provides 17 rotatable bonds of separation between the 3' base of the oligonucleotide and the  $\text{SiO}_2$  surface. To ensure complete amine deprotonation and to minimize secondary structure formation during  
25 coupling, the reaction is performed in 0.1 M KOH and incubated at 37°C for 6 hours.

          In Format 3 SBH in general, signals are scored per each of billion points. It would not be necessary to  
30 hybridize all arrays, e.g., 4000 5 x 5mm, at a time and the successive use of smaller number of arrays is possible.

          Cycling hybridizations are one possible method for  
35 increasing the hybridization signal. In one cycle, most of the fixed probes will hybridize with DNA fragments with tail sequences non-complementary for labelled

- 58 -

probes. By increasing the temperature, those hybrids will be melted (FIG. 3). In the next cycle, some of them (~0.1%) will hybridize with an appropriate DNA fragment and additional labeled probes will be ligated. In this case, there occurs a discriminative melting of DNA hybrids with mismatches for both probe sets simultaneously.

In the cycle hybridization, all components are added before the cycling starts, at the 37°C for T4, or a higher temperature for a thermostable ligase. Then the temperature is decreased to 15-37°C and the chip is incubated for up to 10 minutes, and then the temperature is increased to 37°C or higher for a few minutes and then again reduced. Cycles can be repeated up to 10 times. In one variant, an optimal higher temperature (10-50°C) can be used without cycling and longer ligation reaction can be performed (1-3 hours).

The procedure described herein allows complex chip manufacturing using standard synthesis and precise spotting of oligonucleotides because a relatively small number of oligonucleotides are necessary. For example if all 7-mer oligos are synthesized (16384 probes), lists of 256 million 14-mers can be determined.

One important variant of the invented method is to use more than one differently labeled probe per basic array. This can be executed with two purposes in mind; multiplexing to reduce number of separately hybridized arrays; or to determine a list of even longer oligosequences such as 3 x 6 or 3 x 7. In this case if two labels are used the specificity of the 3 consecutive oligonucleotides can be almost absolute because positive sites must have enough signals of both labels.

- 59 -

A further and additional variant is to use chips containing BxNy probes with y being from 1 to 4. Those chips allow sequence reading in different frames.. This can also be achieved by using appropriate sets of labeled probes or both F and P probes could have some unspecified end positions (i.e., some element of terminal degeneracy). Universal bases may also be employed as part of a linker to join the probes of defined sequence to the solid support. This makes the probe more available to hybridization and makes the construct more stable. If a probe has 5 bases, one may, e.g., use 3 universal bases as a linker (FIG. 4).

15

**EXAMPLE VIII****ANALYZING THE DATA OBTAINED**

Image files are analyzed by an image analysis program, like DOTS program (Drmanac et al., 1993), and scaled and evaluated by statistical functions included, e.g., in SCORES program (Drmanac et al., 1994). From the distribution of the signals an optimal threshold is determined for transforming signal into +/- output.

From the position of the label detected, F + P nucleotide sequences from the fragments would be determined by combining the known sequences of the immobilized and labelled probes corresponding to the labelled positions. The complete nucleic acid sequence or sequence subfragments of the original molecule, such as a human chromosome, would then be assembled from the overlapping F + P sequences determined by computational deduction.

One option is to transform hybridization signals e.g., scores, into +/- output during the sequence assembly process. In this case, assembly will start with

- 60 -

a F+P sequence with a very high score, for example F+P sequence AAAAAATTTTTT (SEQ ID NO:1). Scores of all four possible overlapping probes AAAAAATTTTTTA (SEQ ID NO:3), AAAAAATTTTTTT (SEQ ID NO:4), AAAAAATTTTTTC (SEQ ID NO:5) and AAAAAATTTTTTG (SEQ ID NO:6) and three additional probes that are different at the beginning (TAAAAATTTTTT, SEQ ID NO:7; CAAAAATTTTTT, SEQ ID NO:8; GAAAAATTTTTT, SEQ ID NO:9) are compared and three outcomes defined: (i) only the starting probe and only one of the four overlapping probes have scores that are significantly positive relatively to the other six probes, in this case the AAAAAATTTTTT (SEQ ID NO:1) sequence will be extended for one nucleotides to the right; (ii) no one probe except the starting probe has a significantly positive score, assembly will stop, e.g., the AAAAAATTTTTT (SEQ ID NO:10) sequence is at the end of the DNA molecule that is sequenced; (iii) more than one significantly positive probe among the overlapped and/or other three probes is found; assembly is stopped because of the error or branching (Drmanac et al., 1989).

The processes of computational deduction would employ computer programs using existing algorithms (see, e.g., Pevzner, 1989; Drmanac et al., 1991; Labat and Drmanac, 1993; each incorporated herein by reference).

If, in addition to F + P, F(space 1)P, F(space 2)P, F(space 3)P or F(space 4)P are determined, algorithms will be used to match all data sets to correct potential errors or to solve the situation where there is a branching problem (see, e.g., Drmanac et al., 1989; Bains et al., 1988; each incorporated herein by reference).

- 61 -

## EXAMPLE IX

RE-USING SEQUENCING CHIPS

When ligation is employed in the sequencing process,  
5 then the ordinary oligonucleotides chip cannot be  
immediately reused. The inventor contemplates that this  
may be overcome in various ways.

One may employ ribonucleotides for the second probe,  
10 probe P, so that this probe may subsequently be removed  
by RNAase treatment. RNAase treatment may utilize RNAase  
A an endoribonuclease that specifically attacks single-  
stranded RNA 3' to pyrimidine residues and cleaves the  
phosphate linkage to the adjacent nucleotide. The end  
15 products are pyrimidine 3' phosphates and  
oligonucleotides with terminal pyrimidine 3' phosphates.  
RNAase A works in the absence of cofactors and divalent  
cations.

20 To utilize an RNAase, one would generally incubate  
the chip in any appropriate RNAase-containing buffer, as  
described by Sambrook et al. (1989; incorporated herein  
by reference). The use of 30-50  $\mu$ l of RNAase-containing  
buffer per 8 x 8mm or 9 x 9mm array at 37°C for between  
25 10 and 60 minutes is appropriate. One would then wash  
with hybridization buffer.

Although not widely applicable, one could also use  
the uracil base, as described by Craig et al. (1989),  
30 incorporated herein by reference, in specific  
embodiments. Destruction of the ligated probe  
combination, to yield a re-usable chip, would be achieved  
by digestion with the *E. coli* repair enzyme, uraci-DNA  
glycosylase which removes uracil from DNA.

35

One could also generate a specifically cleavable  
bond between the probes and then cleave the bond after

- 62 -

detection. For example, this may be achieved by chemical ligation as described by Shabarova et al. (1991) and Dolinnaya et al. (1988), both references being specifically incorporated herein by reference.

5

Shabarova et al. (1991) describe the condensation of oligodeoxyribo nucleotides with cyanogen bromide as a condensing agent. In their one step chemical ligation reaction, the oligonucleotides are heated to 97°C, slowly  
10 cooled to 0°C, then 1 µl 10M BrCN in acetonitrile is added.

Dolinnaya et al. (1988) show how to incorporate phosphoramidate and pyrophosphate internucleotide bonds  
15 in DNA duplexes. They also use a chemical ligation method for modification of the sugar phosphate backbone of DNA, with a water-soluble carbodiimide (CDI) as a coupling agent. The selective cleavage of a phosphoamide bonds involves contact with 15% CH<sub>3</sub>COOH for 5 min at  
20 95°C. The selective cleavage of a pyrophosphate bond involves contact with a pyridine-water mixture (9:1) and freshly distilled (CF<sub>3</sub>CO)<sub>2</sub>O.

25

\* \* \*

While the compositions and methods of this invention have been described in terms of preferred embodiments, it  
30 will be apparent to those of skill in the art that variations may be applied to the composition, methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it  
35 will be apparent that certain agents that are both chemically and physiologically related may be substituted for the agents described herein while the same or similar



- 63 -

results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims. All

5 claimed matter and methods can be made and executed without undue experimentation.

## REFERENCES

- The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.
- 10 Bains et al., 1988, J. Theor. Biol., 135:303-307.
- Broude et al., 1994, Proc. Natl. Acad. Sci. USA, 91:3072-3076.
- 15 Brumbaugh et al., 1988, Proc. Natl. Acad. Sci. U.S.A., 85:5610-5614.
- Cantor et al., 1992, Genomics, 13, 1378.
- Cate et al., 1991, GATA, 8(3):102-106.
- 20 Chu et al., 1983, Nucleic Acids Res., 11:6513-6529.
- Craig et al., 1989, Nucleic Acids Research, 17(12):4605.
- 25 Dahlen et al., 1987, Mol. Cell. Probes 1:159-168.
- Dolinnaya et al., 1988, Nucleic Acids Research, 16(9):3721-3738.
- 30 Drmanac & Crkvenjakov, 1990, Scientia Yugoslavica, 16, 97.
- Drmanac & Crkvenjakov, U.S. Patent 5,202,231.
- 35 Drmanac et al., 1989, Genomics, 4:114-128.
- Drmanac et al., 1991, J. Biomol. Struct. & Dyn., 8:1085.
- 40 Drmanac et al., 1991, In "Electrophoreses, Supercomputers and the Human Genome", pp 47-59, World Scientific Publishing Co., Singapore.
- 45 Drmanac et al., 1993, Proceedings of 2nd International Conference on Bioinformatics, Supercomputing, and Complex Genome Analysis, World Scientific Publishing Co., pp. 121-134.
- 50 Drmanac, 1994, Abstract Book for Genome Mapping and Sequencing; arranged by Richard Myers, David Porteous and Robert Waterstone, Cold Spring Harbor Laboratories, p.60.

- 65 -

- Drmanac et al., 1994, Proceedings of the 3rd  
International Workshop of Transcribed Sequences, In  
Press.
- 5 Duncan & Cavalier, 1988, Analytical Biochemistry,  
169:104-108.
- Eggers et al., 1994, BioTechniques, 17(3):516-524.
- 10 Fitzgerald et al., 1992, Nucleic Acids Research,  
20(14):3753-62.
- Fodor et al., 1991, Science, 251:767-768.
- 15 Hoheisel & Lehrach, 1990, FEBS Lett., 274(1,2):103-106.
- Inouye & Hondo, 1990, J. Clin. Microb., 28:1469-1472.
- Jacobsen et al., 1990, Genomics, 8:001-007.
- 20 Keller et al., 1988, Anal. Biochem., 170:441-450.
- Keller et al., 1989, Anal. Biochem., 177:27-32.
- 25 Khrapko et al., 1991, J. DNA Sequencing Mapping, 1, 375.
- Labat and Drmanac, 1993, Proceedings of 2nd International  
Conference on Bioinformatics, Supercomputing, and  
Complex Genome Analysis, World Scientific Publishing  
30 Co., pp. 555-565.
- Landegren et al. 1988, Science, 241:1077-1080.
- Maxam & Gilbert, 1977, Proc. Natl. Acad. Sci., 74, 560.
- 35 Morriey & Collins, 1989, Mol. Cell. Probes 3:189-207.
- Murakami et al., 1991, Nucleic Acids Research,  
19(15):4097-4102.
- 40 Nagata et al., 1985, FEBS Letters, 183:379-382.
- Nichols et al., 1994, Nature, 369:492.
- 45 Pease et al., 1994 Proc. Natl. Acad. Sci., 91:5022-5026.
- Peterkin et al., 1987, BioTechniques 5(2):132-134.
- Peterkin et al., 1989, Food Microbiology 5(2):281-284.
- 50 Pontius & Berg, 1991, Proc. Natl. Acad. Sci. U.S.A.,  
88:8237-8241.
- Rasmussen et al., 1991, Analytical Biochemistry, 198:138-  
55 142.

- 66 -

- Sambrook et al., 1989, Molecular cloning: A laboratory manual. Cold Spring Harbor Laboratory. Cold Spring Harbor, NY.
- 5 Sanger, et al., 1977, Proc. Natl. Acad. Sci., 74, 5463.
- Schriefer et al., 1990, Nucleic Acids Research, 18(24):7455.
- 10 Schubert et al., 1990, Nucleic Acids Research, 18(11):3427.
- Shabarova et al., 1991, Nucleic Acids Research, 19(15):4247-4251.
- 15 Sharp et al., 1989 Food Microbiology, 6:261-265.
- Southern, PCT Patent Application WO 89/10977.
- 20 Southern & Maskos, PCT Patent Application WO 90/03382.
- Southern et al., 1992, Genomics, 13, 1008.
- 25 Strezoska et al., 1991, Proc. Natl. Acad. Sci., 88, 10089.
- Van Ness et al., 1991, Nucleic Acids Research, 19(12):3345.
- 30 Wu & Wallace, 1989 Gene, 76:245-254.
- Zeremski and Crkvenjakow, 1993, DNA Sequence Determination by Hybridization: a Strategy for Efficient Large-Scale Sequencing, Science, 260:1649-35 1652.

- 67 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

5

## (i) APPLICANT:

NAME: ARCH DEVELOPMENT CORPORATION  
STREET: 1101 East 58th Street  
CITY: Chicago  
10 STATE: Illinois  
COUNTRY: United States of America  
POSTAL CODE: 60637  
TELEPHONE NO: (312) 702-1692  
TELEFAX NO: (312) 702-0741

15

(ii) INVENTOR: Drmanac, Radoje

(iii) TITLE OF INVENTION: Methods and Compositions for  
Efficient Nucleic Acid  
20 Sequencing

(iv) NUMBER OF SEQUENCES: 10

## (v) CORRESPONDENCE ADDRESS:

25 (A) ADDRESSEE: Arnold, White & Durkee  
(B) STREET: P. O. Box 4433  
(C) CITY: Houston  
(D) STATE: TX  
(E) COUNTRY: USA  
30 (F) ZIP: 77210-4433

## (vi) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
35 (C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Ver. #1.25

## (vii) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: FILING HEREWITH  
40 (B) FILING DATE: FILING HEREWITH  
(C) CLASSIFICATION:

## (viii) PRIOR APPLICATIONS DATA:

(A) APPLICATION NUMBER: USSN 08/303,058  
45 (B) FILING DATE: 08 SEPTEMBER 1994 (08.09.94)  
(C) CLASSIFICATION: UNKNOWN

(A) APPLICATION NUMBER: USSN 08/127,420  
(B) FILING DATE: 27 SEPTEMBER 1993 (27.09.93)  
50 (C) CLASSIFICATION: UNKNOWN

## (ix) ATTORNEY/AGENT INFORMATION:

(A) NAME: Parker, David L.  
(B) REGISTRATION NUMBER: 32,165  
55 (C) REFERENCE/DOCKET NUMBER: ARCD146P--\PAR

- 68 -

- (x) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: (512) 418-3000  
(B) TELEFAX: (512) 474-7577

5

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 12 base pairs  
10 (B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA  
15

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

AAAAAATTTT TT

12

20

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 13 base pairs  
25 (B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA  
30

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AAAAAATTTT TTC

13

35

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 12 base pairs  
40 (B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: DNA  
45

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AAAAATTTTT TA

12

50

## (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 12 base pairs  
55 (B) TYPE: nucleic acid  
(C) STRANDEDNESS: single

- 69 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AAAAATTTTT TT

12

10 (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

15 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AAAAATTTTT TC

12

25 (2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

30 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

AAAAATTTTT TG

12

40 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

45 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TAAAAATTTTT TT

12

55 (2) INFORMATION FOR SEQ ID NO:8:

- 70 -

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 12 base pairs  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
5      (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:  
10      CAAAAATTTT TT      12

(2) INFORMATION FOR SEQ ID NO:9:  
15      (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 12 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single  
20      (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:  
25      GAAAAATTTT TT      12

(2) INFORMATION FOR SEQ ID NO:10:  
30      (i) SEQUENCE CHARACTERISTICS:  
        (A) LENGTH: 11 base pairs  
        (B) TYPE: nucleic acid  
        (C) STRANDEDNESS: single  
35      (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:  
40      AAAAAATTTT T      11



- 71 -

CLAIMS

1. A method for determining the sequence of a nucleic  
5 acid molecule, comprising the steps of:
  - (a) identifying sequences from the molecule by  
hybridizing the molecule to complementary  
sequences from two sets of small  
10 oligonucleotide probes of known sequence,  
wherein the first set of probes are attached to  
a solid support and the second set of probes  
are labelled probes in solution;
  - 15 (b) identifying overlapping stretches of sequence  
from the sequences identified in step (a); and
  - (c) assembling the nucleic acid sequence of the  
molecule from said overlapping sequences  
20 identified.
2. The method of claim 1, wherein said hybridization is  
carried out in cycles.  
25
3. A method for determining the sequence of a nucleic  
acid molecule, comprising the steps of:
  - 30 (a) fragmenting the nucleic acid molecule to be  
sequenced to provide intermediate length  
nucleic acid fragments;
  - (b) identifying sequences from said fragments by  
35 hybridizing the fragments to complementary  
sequences from two sets of small  
oligonucleotide probes of known sequence,  
wherein the first set of probes are attached to

- 72 -

a solid support and the second set of probes are labelled probes in solution;

5 (c) identifying overlapping stretches of sequence from said sequences identified in step (b); and

(d) assembling the nucleic acid sequence of the molecule from said overlapping sequences identified.

10

4. The method of claim 3, wherein said fragments are sequentially hybridized to complementary sequences from two sets of small oligonucleotide probes of known  
15 sequence.

5. The method of claim 3, wherein said fragments are simultaneously hybridized to complementary sequences from  
20 two sets of small oligonucleotide probes of known sequence.

6. The method of claim 3, wherein said intermediate  
25 length nucleic acid fragments are between about 10 nucleotides and about 40 nucleotides in length and said small oligonucleotide probes are between about 4 nucleotides and about 9 nucleotides in length.

30

7. The method of claim 3, wherein said oligonucleotide probes hybridize to completely complementary sequences from said fragments.

35

- 73 -

8. The method of claim 3, wherein said oligonucleotide probes hybridize to immediately adjacent sequences from said fragments.
- 5 9. The method of claim 8, wherein said oligonucleotide probes hybridize to completely complementary and immediately adjacent sequences from said fragments.
- 10 10. The method of claim 8, wherein said immediately adjacent oligonucleotide probes are subsequently ligated.
11. The method of claim 3, wherein step (b) comprises  
15 the steps of:
- (a) contacting said first set of small attached oligonucleotide probes with said intermediate length nucleic acid fragments under  
20 hybridization conditions effective to allow only those fragments with a completely complementary sequence to hybridize to a probe, thereby forming primary complexes wherein the fragment has hybridized and free sequences;
- 25 (b) contacting said primary complexes with said second set of small labelled oligonucleotide probes under hybridization conditions effective to allow only those probes with completely  
30 complementary sequences to hybridize to a free fragment sequence, thereby forming secondary complexes wherein the fragment is hybridized to an attached probe and a labelled probe;
- 35 (c) removing from said secondary complexes labelled probes that are not immediately adjacent to an

- 74 -

attached probe, thereby leaving only adjacent secondary complexes;

5 (d) detecting said adjacent secondary complexes by detecting the presence of the label; and

(e) identifying sequences from the nucleic acid fragments in said adjacent secondary complexes by connecting the known sequences of the  
10 hybridized attached and labelled probes.

12. A method of nucleic acid sequencing comprising the steps of:

15

(a) fragmenting the nucleic acid to be sequenced to provide nucleic acid fragments of length T;

20

(b) preparing an array of immobilized oligonucleotide probes of known sequences and length F and a set of labelled oligonucleotide probes in solution of known sequences and length P, wherein  $F + P \leq T$ ;

25

(c) contacting said array of immobilized oligonucleotide probes with said nucleic acid fragments under hybridization conditions effective to allow the formation of primary complexes with hybridized, completely  
30 complementary sequences of length F and non-hybridized fragment sequences of length  $T - F$ ;

35

(d) contacting said complexes with said set of labelled oligonucleotide probes under hybridization conditions effective to allow only the formation of secondary complexes with hybridized, completely complementary sequences

- 75 -

of length F and immediately adjacent  
hybridized, completely complementary sequences  
of length P;

- 5 (e) detecting said secondary complexes by detecting  
the presence of the label;
- (f) identifying sequences of length F + P from the  
nucleic acid fragments in said secondary  
10 complexes by combining the known sequences of  
the hybridized immobilized and labelled probes;
- (g) determining stretches of said sequences of  
length F + P that overlap; and
- 15 (h) assembling the complete nucleic acid sequence  
from said overlapping sequences.

20 13. The method of claim 12, wherein length T is about  
three times longer than length F.

14. The method of claim 12, wherein length T is between  
25 about 10 nucleotides and about 40 nucleotides, length F  
is between about 4 nucleotides and about 9 nucleotides  
and length P is between about 4 nucleotides and about  
9 nucleotides.

30 15. The method of claim 14, wherein length T is about  
20 nucleotides, length F is about 6 nucleotides and  
length P is between about 6 nucleotides.

35

- 76 -

16. The method of claim 12, wherein said immediately adjacent immobilized and labeled oligonucleotide probes are ligated.

5

17. A method of nucleic acid sequencing comprising the steps of:

- 10 (a) fragmenting the nucleic acid to be sequenced to provide intermediate length nucleic acid fragments;
- 15 (b) contacting an array of immobilized small oligonucleotide probes of known sequences with said nucleic acid fragments under hybridization conditions effective to allow only those fragments with a completely complementary sequence to hybridize to a probe, thereby forming primary complexes wherein the fragment has hybridized and non-hybridized sequences;
- 20 (c) contacting said primary complexes with a set of labelled small oligonucleotide probes in solution of known sequences under hybridization conditions effective to allow only those probes with completely complementary sequences to hybridize to a non-hybridized fragment sequence, thereby forming secondary complexes wherein the fragment is hybridized to an immobilized probe and a labelled probe;
- 25 (d) removing from said secondary complexes labelled probes that are not immediately adjacent to an immobilized probe, thereby leaving only adjacent secondary complexes;
- 30
- 35

- 77 -

- (e) detecting said adjacent secondary complexes by detecting the presence of the label;
- (f) identifying sequences from the nucleic acid fragments in said adjacent secondary complexes by combining the known sequences of the hybridized immobilized and labelled probes;
- (g) determining stretches of said sequences that overlap; and
- (h) assembling the complete nucleic acid sequence from said overlapping sequences identified.
18. The method of claim 17, wherein the nucleic acid is cloned DNA or chromosomal DNA.
19. The method of claim 17, wherein the nucleic acid is mRNA.
20. The method of claim 17, wherein the nucleic acid is fragmented by restriction enzyme digestion, ultrasound treatment, NaOH treatment or low pressure shearing.
21. The method of claim 17, wherein the nucleic acid fragments are between about 10 nucleotides and about 100 nucleotides in length.
22. The method of claim 17, wherein the oligonucleotide probes are between about 4 nucleotides and about 9 nucleotides in length.

- 78 -

23. The method of claim 22, wherein the oligonucleotide probes are about 6 nucleotides in length.
- 5 24. The method of claim 17, wherein said immobilized oligonucleotides are attached to a glass, polystyrene or teflon solid support.
- 10 25. The method of claim 17, wherein said immobilized oligonucleotides are attached to a solid support via a phosphodiester linkage.
- 15 26. The method of claim 17, wherein said immobilized oligonucleotides are attached to a solid support via a light-activated synthetic mechanism.
- 20 27. The method of claim 17, wherein the labelled oligonucleotide probes are labelled with a non-radioactive isotope or a fluorescent dye.
- 25 28. The method of claim 17, wherein the labelled oligonucleotide probes are labelled with  $^{35}\text{S}$ ,  $^{32}\text{P}$  or  $^{33}\text{P}$ .
29. The method of claim 17, wherein said nucleic acid  
30 fragment or one of said oligonucleotide probes contains a modified base or a universal base.
- 30 30. The method of claim 17, wherein labelled probes that are not immediately adjacent to an immobilized probe are removed from the secondary complexes by stringent washing conditions.



- 79 -

31. The method of claim 17, wherein labelled probes that are immediately adjacent to an immobilized probe are ligated to said immobilized probe and non-ligated. labelled probes are subsequently removed by washing.

5

32. The method of claim 31, wherein said adjacent probes are ligated enzymatically.

10

33. The method of claim 17, wherein multiple arrays of immobilized oligonucleotides are arranged in the form of a sequencing chip.

15

34. A method of nucleic acid sequencing comprising the steps of:

20 (a) fragmenting the nucleic acid to be sequenced to provide nucleic acid fragments of between about 10 nucleotides and about 40 nucleotides in length;

25 (b) contacting an array of immobilized oligonucleotide probes with known sequences of between about 4 nucleotides and about 9 nucleotides in length with said nucleic acid fragments under hybridization conditions effective to allow only those fragments with a completely complementary sequence to hybridize to a probe, thereby forming primary complexes wherein the fragment has hybridized and non-hybridized sequences;

35 (c) contacting said complexes with a set of <sup>32</sup>P-labelled or <sup>33</sup>P-labelled oligonucleotide probes with known sequences of between about

- 80 -

- 4 nucleotides and about 9 nucleotides in length under hybridization conditions effective to allow only those labelled probes with completely complementary sequences to hybridize to a non-hybridized fragment sequence, thereby forming secondary complexes wherein the fragment is hybridized to an immobilized probe and a  $^{32}\text{P}$ -labelled or  $^{33}\text{P}$ -labelled probe;
- 5
- 10 (d) ligating the immobilized probes and labelled probes that are immediately adjacent with a DNA ligase enzyme, thereby forming ligated secondary complexes;
- 15 (e) removing from the secondary complexes any non-ligated labelled probes;
- (f) detecting said ligated secondary complexes by detecting the presence of the  $^{32}\text{P}$  or  $^{33}\text{P}$  label;
- 20
- (g) identifying sequences from the nucleic acid fragments in said ligated secondary complexes by combining the known sequences of the ligated probes;
- 25
- (h) determining stretches of said sequences that overlap; and
- (i) assembling the complete nucleic acid sequence from said overlapping sequences.
- 30

35. A kit for use in nucleic acid sequencing, comprising a solid support chip having attached an arrangement of oligonucleotide probes of known sequences, said oligonucleotides being capable of taking part in hybridization reactions, and a set of containers

35

- 81 -

comprising solutions of labelled oligonucleotide probes of known sequences.

36. The kit of claim 35, wherein multiple chips of  
5 immobilized oligonucleotide probes are arranged in the form of a sequencing array.

37. The kit of claim 35, wherein the oligonucleotide  
10 probes are between about 4 nucleotides and about 9 nucleotides in length.

38. The kit of claim 37, wherein the oligonucleotide  
15 probes are about 6 nucleotides in length.

39. The kit of claim 35, wherein the oligonucleotide  
20 probes are attached to a glass, polystyrene or teflon solid support.

40. The kit of claim 35, wherein the oligonucleotide  
25 probes are attached to a solid support via a phosphodiester linkage.

41. The kit of claim 35, wherein the oligonucleotide  
30 probes are attached to a solid support via a light-activated synthetic mechanism.

42. The kit of claim 35, wherein the labelled  
35 oligonucleotide probes are labelled with a non-radioactive isotope or a fluorescent dye.

- 82 -

43. The kit of claim 35, wherein one of the oligonucleotide probes contains a modified or a universal base.

5

44. The kit of claim 35, wherein the labelled oligonucleotide probes are labelled with  $^{35}\text{S}$ ,  $^{32}\text{P}$  or  $^{33}\text{P}$ .

10 45. The kit of claim 35, further comprising a ligating agent.

15 46. The kit of claim 45, wherein the ligating agent is a DNA ligase enzyme.

1/5




FIG. 1A




FIG. 1B




FIG. 1C

2/5




FIG. 2A




FIG. 2B

3/5




FIG. 2C

4/5




FIG. 3A




FIG. 3B




FIG. 3C



5/5




FIG. 4

# INTERNATIONAL SEARCH REPORT

Internat. J. Application No  
PCT/US 94/10945

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J. DNA SEQUENCE AND MAPPING, vol.1, no.3, 1990, HARWOOD ACADEMIC PUBLISHERS GMBH, UK; pages 375 - 388	1
Y	K.R. KHRAPKO ET AL. 'A method for DNA sequencing by hybridization with oligonucleotide matrix' cited in the application see page 385, left column, line 14 - page 386, left column, line 14; figure 8 --- -/--	2-46

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

10 February 1995

Date of mailing of the international search report

17.02.95

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Hornig, H

## INTERNATIONAL SEARCH REPORT

Internal Application No

PCT/US 94/10945

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GENOMICS, vol.13, no.4, August 1992, ACADEMIC PRESS, NEW YORK, US; pages 1378 - 1383 C.R. CANTOR ET AL. 'Report on the sequencing by hybridization workshop' cited in the application	1
Y	see page 1379, left column, line 10 - line 26; figure 1 ---	2-46
Y	SCIENCE, vol.258, no.5089, 11 December 1992, AAAS, WASHINGTON, DC, US; pages 1787 - 1791 J. KIELECZAWA ET AL. 'DNA sequencing by primer walking with strings of contiguous hexamers' see page 1787, middle column, line 5 - line 16 ---	2-46
Y	GENE, vol.90, no.1, 31 May 1990, ELSEVIER PUBLISHERS, N.Y., U.S.; pages 177 - 178 W. SZYBALSKI 'Proposal for sequencing DNA using ligation of hexamers to generate sequential elongation primers (SPEL-6)' the whole document ---	2-46
Y	WO,A,93 05183 (BAYLOR COLLEGE OF MEDICINE) 18 March 1993 see page 6, line 7 - page 12, line 18; claims 1-26; figure 4 ---	2-46
Y	WO,A,91 06678 (SRI INTERNATIONAL) 16 May 1991 see page 10, line 19 - page 18, line 33; claims 1-50; figure 1 ---	2-46
A	SCIENCE, vol.260, 11 June 1993, AAAS, WASHINGTON, DC, US; pages 1649 - 1652 R. DRMANAC ET AL. 'DNA sequence determination by hybridization: A strategy for efficient large-scale sequencing' the whole document ---	1-46
A	US,A,5 202 231 (DRMANAC ET AL.) 13 April 1993 cited in the application the whole document ---	1-46
	---	
	-/--	

1

# INTERNATIONAL SEARCH REPORT

Intern: A1 Application No  
PCT/US 94/10945

C/(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PROC. NATL. ACAD SCI., vol.86, September 1989, NATL. ACAD SCI., WASHINGTON, DC, US; pages 6917 - 6921 F.W. STUDIER 'A strategy for high-volume sequencing of cosmid DNAs: Random and directed priming with a library of oligonucleotides' the whole document ---</p>	1-46
A	<p>FEBS LETT., vol.256, no.1,2, October 1989, ELSEVIER PUBLISHERS, AMSTERDAM, NL; pages 118 - 122 K.R. KHRAPKO ET AL. 'An oligonucleotide hybridization approach to DNA sequencing' the whole document -----</p>	1-46

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

Intern: Application No

PCT/US 94/10945

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9305183	18-03-93	AU-A- 2674092	05-04-93
WO-A-9106678	16-05-91	EP-A- 0450060	09-10-91
US-A-5202231	13-04-93	NONE	